

Anticipated Classification  
of this Application:  
Class \_\_\_\_\_ Subclass \_\_\_\_\_

06-02-00  
Prior Application:  
Examiner S. McGarry  
Group Art Unit 1635

Attorney's  
Docket  
No. 50995-B/JPW/EMW

June 1, 2000

HONORABLE ASSISTANT COMMISSIONER FOR PATENTS  
Washington, D.C. 20231

06/01/00  
JC822 U.S. PTO  
R:

is a request for filing a X CONTINUATION

DIVISIONAL \_\_\_\_\_ CONTINUATION-IN-PART application under

X 37 C.F.R. §1.53(b) \_\_\_\_\_ 37 C.F.R. §1.53(d), of pending prior application

Serial No. 08/654,482 filed on May 28, 1996 of

Riccardo Dalla-Favera

for

Inventor(s)

IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

Title of Invention

- 06/01/00  
JC822 U.S. PTO  
R:
1. X Enclosed is a copy of the prior application, as originally filed and an affidavit or declaration verifying it as a true copy.
  2. X A verified statement to establish small entity status under 37 C.F.R. §1.9 and 1.27  
\_\_\_\_\_ is enclosed.  
X was filed in the prior application and such status is still proper and desired (37 C.F.R. §1.28(a)).
  3. X The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

	NUMBER FILED		NUMBER EXTRA*		RATE		FEE		
					SMALL ENTITY	OTHER ENTITY		SMALL ENTITY	OTHER ENTITY
Total Claims	7 -20	=	0	X	\$ 9.00	\$18.00	=	\$ 0	\$
Independent Claims	2 -3	=	0	X	\$39.00	\$78.00	=	\$ 0	\$
Multiple Dependent Claims Presented: <u>      </u> Yes <u>  X  </u> No					\$130.00	\$260.00	=	\$ 0	\$
*If the different in Col. 1 is less than zero, enter "0" in Col. 2					BASIC FEE			\$ 345	\$ 690
					TOTAL FEE			\$345.00	\$

4.   X   The Commissioner is hereby authorized to charge payment of the following fees associated with this application or credit any overpayment to Deposit Account No. 03-3125.

  X   Any additional filing fees required under 37 C.F.R. §1.16.

  X   Any patent application processing fees under 37 C.F.R. §1.17.

       The issue fees set forth in 37 C.F.R. §1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. §1.311(b).

5.   X   Three copies of this sheet are enclosed.

6.   X   A check in the amount of \$ 345.00 is enclosed.

7.        Cancel claims                                 .

8.        Amend the specification by inserting before the first line the sentence: --This is a        continuation        division of application Serial No.                         , filed                         .--

9.   X     21   Sheet(s) of        informal   X   formal drawing(s) is/are enclosed.

10.        Transfer the drawings from the prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file.

11.        Priority of application No.                          filed on                          in                          (country) is claimed under 37 U.S.C. §119.

       The certified copy of the priority application has been filed in prior application Serial No.                         , filed                         .

12.   X   The prior application is assigned of record to The Trustees of Columbia University in the City of New York. (a copy of the Assignment is attached)

13.   X   A preliminary amendment is enclosed.

14.   X   The power of attorney in the prior application is to:

John P. White (Reg. No. 28,678); Thomas F. Moran (Reg. No. 16,579); Norman H. Zivin (Reg. No. 25,385); Ivan S. Kavrukov (Reg. No. 25,161); Christopher C. Dunham (Reg. No. 22,031); Thomas G. Carulli (Reg. No. 30,616); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Richard S. Milner (Reg. No. 33,970); Albert Wai-Kit Chan (Reg. No. 36,479); Kristina L. Konstas (Reg. No. 37,864); Mary Anne P. Tanner (Reg. No. 40,197); Timothy X. Witkowski (Reg. No. 40,232); and Mary Catherine DiNunzio (Reg. No. 37,306)

Applicants: Riccardo Dalla-Favera  
U.S. Serial No.: Not Yet Known  
Filed: Herewith  
Cont/Div.  
Page 3

June 1, 2000

- (a)   X   The power appears in the original papers in the prior application.  
(a copy of the Declaration is attached)
- (b)        Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.
- (c)   X   Address all future communications to:  
(May only be completed by applicant,  
or attorney or agent of record.)

John P. White

Cooper & Dunham LLP


1185 Avenue of the Americas

New York, New York 10036

15.   X   Also enclosed Express Mail Certificate of Mailing No. EJ 807 507 669 US, a loose set of loose figures, Statement in Accordance and a computer diskette of Sequence Listing.
16.   X   I hereby verify that the attached papers are a true copy of prior application Serial No. 08/654,482 as originally filed on May 28, 1996.

The undersigned declares further that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statement and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

June 1, 2000  
Date

  
Signature John P. White, Reg. No. 28,678

       INVENTOR(S)  
       ASSIGNEE OF COMPLETE INTEREST  
  X   ATTORNEY OR AGENT OF RECORD  
       FILED UNDER 37 C.F.R. §1.34(a)

Address of Signator:

Cooper & Dunham LLP

1185 Avenue of the Americas

New York, New York 10036

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

Applicant: Riccardo Dalla-Favera  
Serial No.: Not Yet Known (Continuation of U.S. Serial No.  
08/654,482, filed May 28, 1996)  
Filed: Herewith  
For: IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

1185 Avenue of the Americas  
New York, New York 10036  
June 1, 2000

Assistant Commissioner for Patents  
Washington, D.C. 20231  
Box: Patent Application

SIR:

**PRELIMINARY AMENDMENT**

Please amend the above-identified application as follows:

**In the specification:**

At page 1, after the title and before line 3, please insert:

--This application is a continuation of U.S. Serial No. 08/654,482,  
filed May 26, 1996, now allowed, the contents of which are hereby  
incorporated by reference.--

**In the claims:**

Please cancel claims 1-87, 93-97 and 99-101 without prejudice to  
applicant's right to pursue the subject matter of these claims in

Applicant: Riccardo Dalla-Favera  
Serial No.: Not Yet Known(Continuation of U.S. Serial No.  
08/654,482, filed May 28, 1996)  
Filed: Herewith

a future continuation application.

Please amend claims 89-92 and 98 under 37 C.F.R. §1.121(b) by deleting the bracketed materials and inserting the underlined materials as follows:

--89.(Amended) A purified human MUM-1 protein of claim 88[.],wherein the MUM-1 protein has the same amino acid sequence as shown in Figure 5B (SEQ. ID NO:14).--

--90.(Amended) An antibody directed to [a] the purified MUM-1 protein of claim 89.--

--91.(Amended) An antibody capable of specifically recognizing MUM-1 protein, wherein the MUM-1 protein has the same amino acid sequence as shown in Figure 5B (SEQ. ID NO:14).--

--92.(Amended) [An] The antibody of claim 91, wherein the MUM-1 protein is a human MUM-1 protein.--

--98.(Amended) [An] A monoclonal antibody of [any one of claims] claim 90.\_, 91 and 92.]--

Please add new claims 102-103 as follows:

Applicant: Riccardo Dalla-Favera  
Serial No.: 08/654,482  
Filed: May 28, 1996  
Page 3

--102. (New) A monoclonal antibody of claim 91.--

--103. (New) A monoclonal antibody of claim 92.--

#### REMARKS

Claims 1-101 were pending in the subject application. Applicant has hereinabove canceled claims 1-87, 93-97 and 99-101 without disclaimer or prejudice to applicant's right to pursue the subject matter of these claims at a later date. Applicant has also amended claims 89-92 and 98 and added new claims 102-103. Support for amended claims 89 and 91 may be found in the specification inter alia on page 21, lines 2-4 and Figure 5B. Support for amended claim 92 may be found in the specification inter alia on page 32, lines 10-12 and page 21, lines 2-4. Support for amended claim 98 may be found in the specification inter alia on page 32, lines 22-23. Support for new claims 102-103 may be found in the specification inter alia on page 32, lines 23-27 and in previously pending claim 98. Applicant maintains that the amendments are fully supported by the specification and do not raise any issue of new matter. Accordingly, applicant respectfully requests that the Examiner enter the Amendment. Upon entry of the Amendment, claims 88-92 and 98, as amended, and new claims 102-103 will be pending and under examination.


If a telephone interview would be of assistance in advancing prosecution of the subject application, applicant's undersigned attorney invites the Examiner to telephone at the number provided

Applicant: Riccardo Dalla-Favera  
Serial No.: 08/654,482  
Filed: May 28, 1996  
Page 4

below.

No fee is deemed necessary in connection with the filing of this Preliminary Amendment. However, if any additional fee is required, authorization is hereby given to charge the amount of such fee to Deposit Account No. 03-3125.

Respectfully submitted



John P. White  
Registration No. 28,678  
Attorney for the Applicant  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10036  
(212) 278-0400

**Application  
for  
United States Letters Patent**

**To all whom it may concern:**

*Be it known that* I, Riccardo Dalla-Favera

*have invented certain new and useful improvements in*

**IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA**

*of which the following is a full, clear and exact description.*



IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

The invention disclosed herein was made with Government support under NIH Grant No. CA 44025. Accordingly, the U.S. Government has certain rights in this invention.

Background of the Invention

Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains. Full bibliographic citation for these references may be found at the end of this application, preceding the claims.

Multiple myeloma (MM) is an incurable B cell tumor affecting B cell end-stage differentiation. Clinically, the course of MM is similar to end-stage plasma cell leukemia (PCL), i.e., there is an uncontrollable proliferation of myeloma cells accompanied by numerous complications, including hyperviscosity syndromes, hypercalcemia, infections, multiple bone fractures, and organ failure.

Non-random chromosomal translocation is known to play a crucial role in the tumorigenesis of hematologic malignancies (1). In B-cell lymphomas, many important proto-oncogenes deregulated by juxtaposition to immunoglobulin (Ig) gene locus have been identified. Each proto-oncogene is associated with a specific subtype of lymphoma, such as c-MYC in Burkitt's lymphoma, Cyclin D1 in mantle cell lymphoma, BCL-2 in follicular lymphoma and BCL-6 in diffuse large cell lymphoma (2-8). In contrast, little is known about molecular alterations of

human MM/PCL, due to the difficulty in cytogenetic analysis. However, previous cytogenetic reports have shown a 14q+ chromosome, suggesting the existence of a chromosomal translocation involving the Ig heavy chain (IgH) locus, is observed in 20 ~ 30 % of the MM/PCL cases and it is the most frequent consistent abnormality (9-12). Even in such cases, most cytogenetic data have failed to identify donor chromosomes other than 11q13, 8q24, and 18q21, where proto-oncogenes Cyclin D1, BCL-2, and c-MYC are located, respectively. Among them, the 11q13 locus has been demonstrated to be involved in nearly 5~10% of the cases and also in 62% of the established cell lines (13). The t(11;14)(q13;q32) translocation is also accompanied by a corresponding overexpression of the Cyclin D1 gene, which raises a strong possibility of the involvement of this gene, although the breakpoints at 11q13 do not cluster like those of the lymphoma cases (14-16). Recent advances in fluorescence in situ hybridization (FISH) have made it possible to clarify both the frequency of the 14q+ chromosomes and the partner chromosomes of the IgH loci. One such report revealed an intriguing result, i.e., that numerous chromosomal loci are able to translocate to IgH locus, including 6p21, 1q21, 3p11, 7q11, 11q23 (17). This has prompted a search for the proto-oncogenes deregulated by the regulatory elements of the IgH gene for a further understanding of the molecular mechanisms of MM/PCL. In the present study, one candidate proto-oncogene, MUM1 (multiple myeloma oncogene 1), was found juxtaposed to the IgH gene as a result of t(6;14)(p25; q32) translocation in human myeloma cell line, SKMM-1. Over expression of the MUM1 mRNA was observed in this cell line. A second gene, called MUM-2 was found translocated in proximity to the IgH gene on

chromosome 14q32 in human myeloma cell line, U-266.

The method of analysis of 14q+ chromosomal translocations and identification of the genes altered in multiple myeloma of this invention are useful since 1) no method is currently available to determine the chromosomal sequences involved in 14q+ translocations, the most important cytogenetic lesions associated with MM pathogenesis; 2) no specific gene lesion is currently known for MM; 3) no diagnostic method based on gene/DNA lesion is currently available for MM and 4) there are no therapeutic approaches aimed at counteracting the action of abnormal gene products in MM.

### Summary of the Invention

This invention provides a method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of: (a) obtaining a DNA sample from the subject suffering from multiple myeloma; (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample; (c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction; (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene; (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones from step (d); (f) hybridizing said fluorescent probes with metaphase chromosomes; and (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma.

This invention provides a method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma which comprises steps of: a) selecting a probe having a sequence of a chromosome other than chromosome 14, identified at the chromosomal breakpoint detected in a subject suffering from multiple myeloma, wherein said probe is capable of hybridizing to the unique sequence of the gene other than

the immunoglobulin gene altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma; b) contacting said probe with mRNA isolated from a cell under conditions permitting formation of a complex between said probe and the mRNA; c) isolating the complex resulting from step (b); d) determining the sequence of the mRNA in the isolated complex, thereby determining the identity of the gene.

10 This invention provides a gene designated *MUM-1*. This invention provides a gene designated *MUM-2*. This invention provides an isolated nucleic acid molecule encoding a MUM protein. This invention provides a DNA encoding a MUM protein. This invention provides a cDNA encoding a MUM protein. This invention provides a genomic DNA molecule encoding a MUM protein. This invention provides a RNA molecule encoding a MUM protein. This invention provides an isolated nucleic acid molecule encoding a human *MUM-1* protein. This invention provides an isolated nucleic acid molecule encoding a human *MUM-2* protein. This invention provides an isolated nucleic acid molecule encoding a MUM protein operatively linked to a promoter of RNA transcription. This invention provides a vector comprising the an isolated cDNA encoding a MUM protein. This invention provides a vector which comprises an isolated cDNA encoding a MUM protein. This invention provides a vector which comprises an isolated cDNA encoding a MUM protein, wherein the vector is a plasmid. This invention provides a host cell for the vector which comprises an isolated cDNA encoding a MUM protein.

This invention provides a nucleic acid probe comprising a

nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a MUM protein. This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM protein.

This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

This invention provides a nucleic acid probe comprising a the sequence of a nucleic acid molecule encoding a MUM-1 protein which is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

This invention provides a nucleic acid probe comprising a the sequence of a nucleic acid molecule encoding a MUM-2 protein which is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein. This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the

subject a rearrangement of nucleic acid encoding MUM-2 protein.

5 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule  
10 encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule.

15 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an isolated cDNA molecule encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the isolated genomic DNA molecule encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence  
20 capable of specifically hybridizing to an isolated RNA molecule encoding a MUM protein.

25 This invention provides a purified MUM protein. This invention provides a purified MUM-1 protein. This invention provides an antibody directed to a purified MUM-1 protein. This invention provides an antibody capable of specifically recognizing MUM-1 protein. This invention provides a purified MUM-2 protein. This invention provides an antibody directed to a purified MUM-2 protein. This invention  
30 provides an antibody capable of specifically recognizing a MUM-2 protein.

This invention provides a pharmaceutical composition

comprising an amount of an oligonucleotide effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane. This invention provides a

5 pharmaceutical composition comprising an amount of the oligonucleotide effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.



**Brief Description of the Figures**

Figure 1. JH-C $\mu$  dissociation in *Bam*HI digested DNA of the 14q+ SK-MM-1 cell line. A 10  $\mu$ g of the high molecular weight DNA was completely digested with *Bam*HI, loaded on each lane and blotted. The same filter was sequentially hybridized with JH, C $\mu$ , Cy2, and 0.7B/H probes. JH probe detects two rearranged bands of 12.0 kb and 9.7kb. The 9.7 kb band is comigrated with that probed with Cy2 probe, suggesting it to be a physiological rearrangement. On the other hand, one allele of the C $\mu$  locus is deleted and another is rearranged (6.5 kb) without being comigrated with rearranged bands of JH. Therefore, 12.0 kb and 6.5 kb bands detected by JH and C $\mu$  (shown by arrowheads) might represent unknown derivative chromosome and derivative 14 chromosome, respectively. As expected, 0.7B/H probe (Fig. 2A) detected the rearranged band comigrated with 6.5kb band of C $\mu$ . Dashed lines show the comigration. Size markers of  $\lambda$ /*Hind*III are shown on the left.

Figures 2A-B. Molecular cloning of the breakpoints of the t(6;14) translocation and germline walking at MUM1 locus. (A) Restriction maps of  $\lambda$ SKB-4a and  $\lambda$ SKS-3 clones representing derivative 6 and 14 are shown, together with germline maps of IgH locus at 14q32 and MUM1

locus at 6p25. Arrows indicate the chromosomal breakpoints. B, *Bam*HI; E, *Eco*RI; H, *Hind*III. (B) Comparison of the nucleotide sequences around the breakpoints on derivative 6 and derivative 14 chromosome. Homologous regions are indicated by dashes. The arrow indicates the breakpoint. Nucleotide numbers shown below are the same as in the Sp sequence reported by Sun, et al. (18).

Figure 3. Mapping of the MUM1 locus to chromosome 6p25.  $\lambda$ MUM-3 genomic clone (Figure 2A) was used as a probe for *in situ* hybridization. The white arrow indicates the fluorescence signal on chromosome 6 band p25. Right panel shows the G-banding picture stained with DAPI.

Figures 4A-C. Expression of the MUM1 gene in hematopoietic lineage. A 10 ug aliquot of total RNA was loaded on each lane and Northern blot analysis was performed using the 2.1H probe (Figure 2A). GAPDH or  $\beta$ -actin probes were used to control for amount of RNA loaded. (A) MUM1 RNA expression in various hematopoietic cell lines. MUM1 RNA is detected in B cell and mature T cell lines as a single 6kb transcript. HELA, epithelial lineage; LCL, Epstein-Barr virus-transformed lymphoblastoid cell line; RAMOS and SK-MM-1, B-cell lineage; HUT-78 and MOLT-4, T-cell

lineage; HL-60 and U937, myelomonocytic lineage; K562, erythroid lineage. Dashes indicate 28S and 18S. (B) Expression in B cell lines derived from various stages of B cell differentiation. MUM1 RNA is seen throughout the B cell development except for BJAB cell line. 697, pre-B cell stage; RAMOS and BJA-B, Burkitt cell line representing mature-B cell stage; RPMI-8226 and U-266, plasma cell stage. (C) Comparison of the expression level among myeloma cell lines. MUM1 RNA is overexpressed in SK-MM-1 cell line carrying t(6;14). Overexpression of the MUM1 is also demonstrated in XG-4, XG-7, and XG-10 cell lines. RPMI-8226, U-266, EJM, and SKMM-1 are IL-6 (interleukin-6) independent lines, whereas XG-1, XG-2, XG-4, XG-5, XG-6, XG-7, and XG-10 are IL-6 dependent lines.

Figures 5A-B. Sequence of MUM1 cDNA and structure of its predicted protein product. (A) Restriction map of the MUM1 cDNA and the position of the open reading frame (box). The solid box indicates approximate position of the DNA binding domain. Sc, *SacII*; A, *ApaI*; P, *PstI*; H, *HindIII*; S, *SacI* (B) Nucleotide sequence of the MUM1 cDNA and corresponding amino acid sequence. Putative translation initiation codons and preceding stop codons appearing in frame are underlined. The asterisk indicates the translation stop

codon.

Figures 6A-B. Homology between MUM1 and other IRF family proteins. (A) Similarity at N-terminal DNA binding domain. Black background indicates identical residues found more than four times. Gray indicates conserved residues that appear in at least four sequences at a given position. Conserved tryptophan residues in DNA binding domain among IRF family members are indicated by closed circles. (B) Similarity at C-terminal region between human MUM1, Mouse LSIRF/Pip, Human ICSBP, Human ISGF3 $\gamma$ , and Human IRF-3. Black and gray background are as in (A).

Figure 7. Genomic organization of the MUM1 gene and location of the chromosomal breakpoints in multiple myeloma. Filled boxes indicate the coding regions and empty boxes indicate the noncoding regions. The position, and the size of each exon of the MUM1 gene are approximate and have been determined by the hybridizations. One exon in each restriction fragment may consist of more than two exons. Translation initiation codon (ATG) and stop codon (TGA) are indicated. Genomic probes used for further investigations are shown as solid bars below the map. Arrows indicate the chromosomal breakpoints of SKMM-1 cell line and case 10. B, *Bam*HI; E, *Eco*RI; H, *Hind*III.

Figure 8. Scheme of the t(6;14)(p25;q32) translocation involving the MUM1 and the immunoglobulin heavy chain (IgH) gene loci. VH-D-J-CH indicates variable-diversity joining-constant region of the IgH gene. Direction of the MUM1 gene on the chromosome 6 is tentatively drawn.

Figure 9A-B. Demonstration of JH-Cα disjunction in U-266 cells and cloning of normal and 14q+ chromosomal breakpoints. (A) The panel shows the results of Southern blot analysis of BamHI digested U-266 and normal control (placenta) genomic DNA using the indicated JH and Cα probes. The arrowheads indicate two DNA fragments containing Cα sequences not linked to JH sequences, suggesting the presence of a chromosomal breakpoint in 14q32. (B) The panel provides a schematic representation of the phage clones isolated from a library constructed from U-266 DNA and screened with a Cα probe. Based on restriction enzyme analysis, the three cloned regions represent a normal Cα region (14q32 germ-line), and two rearranged regions (der.14 and 14q32) containing unknown sequences linked to Cα sequences. The 2.5BE probe used for Northern blot analysis of MUM2 transcripts (Fig. 10) is also shown.

Figure 10. Identification of MUM2 RNA transcripts. The

figure shows the results of a Northern blot analysis of RNA extracted from various MM/PCL cell lines using the 2.5BE probe (see Fig. 9) or GAPDH probe (as a control for RNA loading). A 1.9 Kb RNA transcript is detectable in some cell lines including U-266, indicating that the 2.5BE fragments represents part of a gene, MUM2.

10 Figure 11A-B. Schematic representation of IgH DNA rearrangements in normal B cells and in tumors carrying chromosomal translocations breaking the S region of the IgH locus. Note that in physiological IgH rearrangements (panel 11A) JH sequences and C sequences (C<sub>μ</sub> before and C<sub>γ</sub> after switch recombination, respectively) are consistently found within the same BamHI restriction fragment. Conversely, JH and C sequences are not linked, and are present on two different chromosomes [derivative X and derivative 14(14q+)] in cells carrying a chromosomal translocation breaking the switch region (panel 11B)

25 Figure 12A-B. MUM1 cDNA. cDNA insert is cloned into EcoRI/BamHI site of the pBluescript KS+. Bacteria strain used is DH5α cells. pcMUM1.16a contains full length open reading frame of nt. 217-1572.

30 Figure 13. Breakpoint Cloning of the U-266 Cell Line.

-15-

pMUM2-8 has a 22.0 KB insert in BamHI site  
of pBluescript KS+.

**Detailed Description of the Invention**

The following standard abbreviations are used throughout the specification to indicate specific nucleotides:

5

C=cytosine      A=adenosine  
T=thymidine    G=guanosine

10

This invention provides a method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of: (a) obtaining a DNA sample from the subject suffering from multiple myeloma; (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample;

15

(c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction; (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene; (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones

20

from step (d); (f) hybridizing said fluorescent probes with metaphase chromosomes; and (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma.

25

30

In an embodiment, step (b) of the above described method of this invention is performed by Southern blotting. In another embodiment, step (b) of the above method of this



invention is performed by polymerase chain reaction (PCR) with appropriate probes. Polymerase chain reaction is well known in the art. Since the sequences of both the C and J regions of an immunoglobulin heavy chain gene are known,  
5 appropriate probes for PCR may routinely be designed.

In an embodiment, the genomic library is a phage vector library. In another embodiment, the genomic DNA fragments are generated by cleaving genomic DNA from cells of the  
10 subject with an appropriate restriction enzyme. In a further embodiment, the restriction enzyme is *Bam*HI. In an embodiment, the restriction enzyme is *Sau*3AI. In another embodiment, the probe of step (d) is a human IgH J region JH probe. In a further embodiment, the probe of step (d) is a  
15 human IgH C $\mu$  probe. In an embodiment, the probe of step (d) is a human IgH C $\gamma$ 2 probe. In another embodiment, the chromosomal breakpoint identified is a t(6;14)(p25;q32) translocation. In an embodiment, the chromosomal breakpoint identified is a t(14;15) translocation.

This invention provides a method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma which comprises  
20 steps of: a) selecting a probe having a sequence of a chromosome other than chromosome 14, identified at the chromosomal breakpoint detected in a subject suffering from multiple myeloma, wherein said probe is capable of hybridizing to the unique sequence of the gene other than  
25 the immunoglobulin gene altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma; b) contacting said probe with mRNA isolated from a cell under conditions permitting formation of a complex between said probe and the mRNA; c) isolating the complex resulting from  
30 step (b); and d) determining the sequence of the mRNA in the  
35

isolated complex, thereby determining the identity of the gene.

5 In an embodiment, step (d) of the method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma comprises steps of: i) synthesizing complementary DNA to the mRNA; and ii) performing sequence analysis of the complementary DNA to  
10 determine the sequence of the mRNA.

15 This invention provides a gene identified by the method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma.

20 As used herein, "MUM" means any gene rearranged in 14q+ chromosomal abnormalities associated with multiple myeloma.

This invention provides a gene identified by the above method designated *MUM-1*. This invention provides a gene identified by the above method designated *MUM-2*.

25 This invention provides a gene identified by the above method, wherein the gene identified comprises a nucleic acid encoding a MUM protein. In an embodiment, the gene identified by the above method comprises a nucleic acid encoding a MUM-1 protein. In another embodiment, the gene  
30 identified by the above method comprises a nucleic acid encoding a MUM-2 protein.

This invention provides an isolated nucleic acid molecule encoding a MUM protein. In an embodiment, the isolated

nucleic acid molecule encoding a MUM protein is a DNA molecule. In another embodiment, the isolated nucleic acid molecule encoding a MUM protein is a cDNA molecule.

5

In an embodiment, a cDNA nucleic acid molecule encoding a MUM-1 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as pcMUM1-1.6a (ATCC Accession No. \_\_\_\_\_). Plasmid pcMUM1-1.6a was deposited on  
10 May 28, 1996 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. Plasmid pcMUM1-1.6a  
15 was accorded ATCC Accession Number \_\_\_\_\_.

In another embodiment, a partial cDNA nucleic acid molecule encoding a MUM-1 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as pMUM1-2.4B/N  
20 (ATCC Accession No. \_\_\_\_\_). Plasmid pMUM1-2.4B/N was deposited on May 28, 1996 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of  
25 Microorganisms for the Purposes of Patent Procedure. Plasmid pMUM1-2.4B/N was accorded ATCC Accession Number  
\_\_\_\_\_.

In another embodiment, a partial cDNA nucleic acid molecule  
30 encoding a MUM-1 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as pMUM1-7.7B (ATCC Accession No. \_\_\_\_\_). Plasmid pMUM1-7.7B was deposited on May 28, 1996 with the American Type Culture Collection

(ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. Plasmid pMUM1-7.7B  
5 was accorded ATCC Accession Number \_\_\_\_\_.

In another embodiment, a partial cDNA of the nucleic acid molecule encoding a MUM-2 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as  
10 pMUM2-8 (ATCC Accession No. \_\_\_\_\_). Plasmid pMUM2-8 was deposited on May 28, 1996 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of  
15 Microorganisms for the Purposes of Patent Procedure. Plasmid pMUM2-8 was accorded ATCC Accession Number \_\_\_\_\_.

In an embodiment, the isolated DNA molecule encoding a MUM protein is a cDNA molecule having the nucleotide sequence  
20 shown in Figure 5B (SEQ. ID NO \_\_\_\_\_).

In an embodiment, the isolated DNA molecule encoding a MUM protein is genomic DNA molecule. In an embodiment, the isolated nucleic acid molecule encoding a MUM protein is an  
25 RNA molecule.

In an embodiment, the isolated nucleic acid encodes a human MUM-1 protein. In another embodiment, the isolated nucleic acid molecule encodes a human MUM-2 protein.

30 In an embodiment, isolated nucleic molecule encodes the a human MUM-1 protein having substantially the same amino acid

sequence as shown in Figure 5B (SEQ. ID NO ). In an embodiment, isolated nucleic molecule encodes a human MUM-1 protein having the same amino acid sequence as shown in Figure 5B (SEQ. ID NO ). In an embodiment, the isolated  
5 nucleic acid molecule encoding a MUM protein is operatively linked to a promoter of RNA transcription.

This invention provides a vector comprising a cDNA molecule encoding a MUM protein. In an embodiment, a vector  
10 comprising cDNA encoding for MUM-1 is designated pcMUM1.6a. In an embodiment, a vector comprising partial cDNA encoding for MUM-1 is designated pMUM1.2.4B/N. In an embodiment, a vector comprising partial cDNA encoding for MUM-1 is designated pMUM1-7.7B. In an embodiment, a vector  
15 comprising partial cDNA encoding for MUM-2 is designated pMUM2-8. In an embodiment, a vector comprises genomic DNA encoding for MUM. In an embodiment, the vector is a plasmid. In an embodiment, a host cell comprises the vector comprising cDNA encoding for MUM. In an embodiment, a host  
20 cell comprises the vector comprising genomic DNA encoding for MUM. In a further embodiment, the host cell comprising vectors comprising cDNA encoding for MUM or comprising genomic DNA encoding for MUM is selected from a group consisting of a bacterial cell, a plant cell, and insect  
25 cell and a mammalian cell.

This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a unique sequence included  
30 within the sequence of a nucleic acid molecule encoding a MUM protein. This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the

isolated nucleic acid molecule encoding a MUM protein.

As used herein, the phrase "specifically hybridizing" means the ability of a nucleic acid molecule to recognize a nucleic acid sequence complementary to its own and to form double-helical segments through hydrogen bonding between complementary base pairs.

In an embodiment, the nucleic acid probe specifically hybridizes with nucleic acid encoding MUM-1. In an embodiment, the nucleic acid probe is complementary to nucleic acid encoding MUM-1. In an embodiment, the nucleic acid probe specifically hybridizes with nucleic acid encoding MUM-2. In an embodiment, the nucleic acid probe is complementary to nucleic acid encoding MUM-2.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is a DNA probe. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is a DNA probe.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is a RNA probe. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is a RNA probe.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is a genomic DNA probe. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is a genomic DNA probe.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is labeled with

a detectable marker. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is labeled with a detectable marker.

5 In an embodiment, the detectable marker is selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

10 In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is linked to a nucleic acid sequence capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule of human chromosome 14. In an embodiment, the nucleic acid probe which specifically  
15 hybridizes with nucleic acid encoding MUM-2 is linked to a nucleic acid sequence capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule of human chromosome 14.

20 This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid  
25 molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid  
30 molecule encoding a MUM-1 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM-2 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM-1 protein which is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM-2 protein which is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the specific break point of the nucleic acid probe comprises a portion of the t(6;14)(p25;q32) translocation. In an embodiment, the specific break point of the nucleic acid probe comprises a portion of a t(14;15) translocation. In an embodiment, the nucleic acid probe comprising a portion of the t(6;14)(p25;q32) translocation is labeled with a detectable marker. In an embodiment, the nucleic acid probe comprising a portion of a t(14;15) translocation is labeled with a detectable marker. In an



embodiment, the nucleic acid probe comprising a portion of the t(6;14)(p25;q32) or comprising a portion of a t(14;15) translocation of claim 60, has a detectable marker selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein.

This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein.

In an embodiment, the rearrangement of nucleic acid encoding MUM-1 protein is detected by contacting the nucleic acid from the sample with a MUM-1 probe under conditions permitting the MUM-1 probe to hybridize with the nucleic acid encoding MUM-1 protein from the sample, thereby detecting the rearrangement of nucleic acid encoding MUM-1 protein in the sample.

In an embodiment, the rearrangement of nucleic acid encoding MUM-2 protein is detected by contacting the nucleic acid from the sample with a MUM-2 probe under conditions permitting the MUM-2 probe to hybridize with the nucleic acid encoding MUM-2 protein from the sample, thereby

detecting the rearrangement of nucleic acid encoding MUM-2 protein in the sample.

5 In an embodiment, the rearrangement of nucleic acid encoding MUM-1 protein is detected by a MUM-1 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

15 In an embodiment, the rearrangement of nucleic acid encoding MUM-2 protein is detected by a the MUM-2 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

20 In an embodiment, the MUM-1 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

30 In an embodiment, the MUM-2 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

35 In an embodiment, the MUM-1 probe comprises a specific break point comprising a portion of the t(6;14)(p25;q32)

translocation. In an embodiment, the MUM-2 probe comprises a specific break point comprising a portion of a t(14;15) translocation.

In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein comprises: a) obtaining DNA from the sample of the subject suffering from multiple myeloma; b) performing a restriction digest of the DNA with a panel of restriction enzymes; c) separating the resulting DNA fragments by size fractionation; d) contacting the resulting DNA fragments with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker; e) detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma; f) preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and g) comparing the detected band pattern specific to the DNA obtained from a sample of subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns

are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

5 In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein  
10 comprises: a) obtaining DNA from the sample of the subject suffering from multiple myeloma; b) performing a restriction digest of the DNA with a panel of restriction enzymes; c) separating the resulting DNA fragments by size fractionation; d) contacting the resulting DNA fragments  
15 with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point  
20 to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker; e) detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule  
25 encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple  
30 myeloma; f) preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and g) comparing the detected band pattern specific to the DNA obtained from a sample of subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for

diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

5

In an embodiment, the size fractionation in step (c) is effected by a polyacrylamide or agarose gel. In an embodiment, the detectable marker is radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

10

In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein comprises: a) obtaining RNA from the sample of the subject suffering from multiple myeloma; b) separating the RNA sample by size fractionation; c) contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker; d) detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma; e) preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and f) comparing the detected band pattern specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from

15

20

25

30

35

step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

5 In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein  
10 comprises: a) obtaining RNA from the sample of the subject suffering from multiple myeloma; b) separating the RNA sample by size fractionation; c) contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included  
15 within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 15 and labeled with a detectable marker; d)  
20 detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma; e) preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and f) comparing the detected band pattern  
25 specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple  
30 myeloma if the patterns are the same.

In an embodiment, the size fractionation in step (b) is effected by a polyacrylamide or agarose gel. In an embodiment, the detectable marker is radioactive isotope,

enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

5 In an embodiment, multiple myeloma associated with the expression of a specific human MUM-1 is diagnosed by the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a DNA or RNA sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid  
10 encoding MUM-1 protein.

15 In an embodiment, multiple myeloma associated with the expression of a specific human MUM-2 is diagnosed by the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-12 protein in a DNA or RNA sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein.

20 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule. This invention provides an antisense oligonucleotide having a sequence  
25 capable of specifically hybridizing to an mRNA molecule encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule.

30 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the cDNA molecule encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the genomic DNA molecule

encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the RNA molecule encoding a MUM protein.

5 This invention provides a purified MUM protein. This invention provides a purified MUM-1 protein. This invention provides a purified human MUM-1 protein. This invention provides an antibody directed to a purified MUM-1 protein. This invention provides an antibody capable of specifically  
10 recognizing MUM-1 protein. In an embodiment, the antibody capable of specifically recognizing MUM-1 protein is a human MUM-1 protein.

This invention provides a purified MUM-2 protein. This  
15 invention provides a purified human MUM-2 protein. This invention provides an antibody directed to a purified MUM-2 protein. This invention provides an antibody capable of specifically recognizing MUM-2 protein. In an embodiment, the antibody capable of specifically recognizing MUM-2  
20 protein is a human MUM-2 protein.

In an embodiment, the antibody directed to a purified MUM-1 protein is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-1 protein  
25 is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-1 protein is a human MUM-1 protein.

In an embodiment, the antibody directed to a purified MUM-2  
30 protein is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-2 protein is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-2 protein is a human



MUM-2 protein.

5 This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier capable of passing  
10 through a cell membrane.

15 This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to a cDNA molecule encoding a MUM protein effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

20 This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to a genomic DNA molecule effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier  
25 capable of passing through a cell membrane.

30 This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing

through a cell membrane.

5 This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to a cDNA molecule encoding a MUM protein effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

10

This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to a genomic DNA molecule effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

15

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

20

25

## **EXPERIMENTAL DETAILS**

### **Materials and Methods**

5     **Cell lines.** The following myeloma cell lines were used in  
the present study: SK-MM-1, RPMI-8226, U266, EJM, XG-1,  
XG-2, XG-4, XG-5, XG-6, XG-7, and XG-10. The RPMI-8226 cell  
line was obtained through the American Type Culture  
Collection (ATCC, Rockville, MD). SKMM-1 and U-266 cell  
10    lines were gifts from Dr. A. N. Houghton and Dr. K. Nilsson,  
respectively (18; 12). Characterization of these cell lines  
were previously reported. Six XG cell lines were gifts from  
Dr. B. Klein and were cultured in RPMI 1640 containing 10%  
fetal calf serum (FCS),  $5 \times 10^{-6}$  mol/L 2-ME, and rIL-  
15    6 (1ng/mL) (13;19). Other myeloma cell lines used were all  
IL-6 independent. The SK-MM-1 cell line was used to isolate  
the chromosomal breakpoint carrying the 14q+ chromosome  
without any information on the donor chromosome. XG-1,  
XG-2, XG-6, XG-8 cell lines are reported to carry the  
20    t(11;14)(q13;q32) translocation. XG-5 cells also share  
both t(11;14) and t(8;14)(q24;q32).

**Southern and Northern blot analyses.** Southern blot analysis  
was performed as previously described (21). Briefly, ten  
25    micrograms of high molecular-weight DNA extracted from each  
cell line was digested to completion with *Bam*HI and *Hind*III  
restriction enzymes, size- fractionated on 0.7% agarose gel,  
and transferred onto Duralose nitrocellulose membrane  
(Stratagene) according to the manufacturer's instructions.  
30    Blots were hybridized with a random-primed DNA probe and  
washed at 60°C in 0.2 x SSC and 0.1 % SDS for 5 minutes.  
Genomic probes used in this study were as follows; human IgH

J region JH probe (6.6kb *Bam*HI-*Hind*III fragment) was provided by Dr. J. V. Ravetch, human IgH C $\mu$  probe (1.3 kb *Eco*RI fragment) was provided by Dr. S.J. Korsmeyer. Human IgH region Cy2 probe was provided by Dr. C. Croce.

5

Northern blot analysis was performed as described previously (21). Briefly, a 10  $\mu$ g aliquot of total RNA was loaded on each lane and probed with a 2.1H probe of the MUM1 gene (Figure 2A). GAPDH or  $\beta$ -actin probes were used as controls for amount of total RNA.

10

**Genomic library.** High molecular-weight DNA of SK-MM-1 cell line was digested completely with *Bam*HI and partially with *Sau*3AI, and size-fractionated by using a low-melting point agarose gel. DNA ranging from 10kb to 23kb were purified and ligated into the *Bam*HI sites of  $\lambda$ -DASH II phage vector (Stratagene, La Jolla, CA). After packaging,  $3 \times 10^5$  and  $6 \times 10^5$  recombinant clones of the *Bam*HI digested library and partially digested library were screened with JH and C $\mu$  probes, respectively. To isolate the germline region of the 6p25 locus, a commercially available human placental library (Stratagene) was screened. Positive clones were mapped with restriction enzymes by partial digestion of the phage DNAs followed by probing with T7 and T3 primers labeled with T4 polynucleokinase and 32p- $\gamma$ ATP.

15

20

25

**cDNA library.** A phage library constructed by oligo-dT and random-priming normal human spleen RNA (Clontech) was screened by 2.1H probe (Figure 2A) to isolate initial MUM1 cDNA clones. After the first round of screening, positive clones were used as probes to walk to the 5' side using the same library. Positive clones were subcloned into

30

pBluescript and analyzed for mapping and sequencing.

**DNA sequencing.** DNA sequences were determined by the dideoxy chain termination method and analyzed by an  
5 ABI (Applied Biosystems) autosequencer. Deletion mutants for sequencing were prepared using exonuclease III and mung bean nuclease. cDNA sequences were analyzed with the Genetics Computer Group (GCG) programs. Sequence homology searches  
10 were carried out through the BLAST E-mail server at the National Center for Biotechnology Information, National Library of Medicine, Bethesda, MD.

**Fluorescence in situ hybridization (FISH).** Metaphase chromosome from human lymphocytes were prepared. A  
15 biotin-labeled probe was prepared by nick-translation using Bio-16-dUTP. Conditions for hybridization and washing were described previously (22).

### **Experimental Results**

20 **IgH gene rearrangement of the SK-MM-I cell line.** In *Bam*HI digestion, the JH probe detects two rearranged bands of the size of 12.0 kb and 9.7kb (Fig 1). The 9.7 kb band is comigrated with that probed with C $\gamma$ 2 probe, suggesting it to  
25 be a physiological rearrangement, although this cell line secretes only  $\lambda$  chain. One allele of the C $\mu$  locus is deleted and another is rearranged (6.5 kb) without being comigrated with rearranged bands of JH. Hybridization with a C $\alpha$  probe showed only the germline band (data not shown).  
30 These results suggested the possibility of the chromosomal breakpoint between JH and C $\mu$  locus. Hence, the 12.0 kb and the 6.5 kb bands detected by JH and C $\mu$  were considered to

represent unknown derivative chromosome and derivative 14 chromosome, respectively.

**Molecular cloning of the t(6;14)(p25;q32) breakpoint.**

5 genomic library constructed with *Bam*HI complete digestion was screened with a JH probe to isolate the 12.0 kb *Bam*HI band. Another library constructed with *Sau*3AI partial digestion was screened with a C $\mu$  probe to isolate phage clones containing the 6.5 kb *Bam*HI fragment. Two phage clones,  $\lambda$ SKB-4a and  $\lambda$ SKS-3, considered to represent the unknown derivative and derivative 14 chromosomes respectively, were obtained (Fig 2A). A 0.7 kb *Bam*HI-*Hind*III probe (0.7B/H) of the  $\lambda$ SKS-3 was used to confirm the comigration with the rearranged 6.5 kb C $\mu$  band by Southern analysis (Fig 1). The chromosomal origin of the centromeric side of the  $\lambda$ SKB-4a and telomeric side of the  $\lambda$ SKS-3 were confirmed by hybridization to a somatic cell hybrid DNA panel with a 4.5 kb *Apa*I fragment (4.5A) and 2.1 kb *Hind*III (2.1H) probes. Both probes showed positive signals in hybrid cell DNA containing a human chromosome 6 (data not shown). These probes were also used to isolate the germline chromosome 6 region by screening the human placental genomic library. One of the phage clone DNA ( $\lambda$ MUM-3) was used as a probe for FISH analysis. It identified the localization of this region to be chromosome 6 short arm p25 (Fig 3). To investigate the precise breakpoint within the IgH gene, a 1.5 kb *Hind*III-*Eco*RI fragment of the  $\lambda$ SKS-3, containing the breakpoint on derivative 14 chromosome was sequenced. The breakpoint was confirmed to be just 3' to the switch  $\mu$ (S  $\mu$ ) repetitive sequences (Fig 2B). Nucleotide sequencing of the region around the breakpoints of chromosome 6 and derivative 6

chromosome showed that the chromosomal translocation was reciprocal with minimum deletion of both the IgH and 6p25 sequences.

5     **Transcriptional unit in the vicinity of the 6p25 breakpoint.**

10     An attempt to find a functional transcriptional unit in the vicinity of the breakpoints was made. Although a 4.5A probe on derivative 6 chromosome could not detect any transcripts, a 2.1H probe on derivative 14 chromosome detected a single  
15     6 kb transcript in the SK-MM-1 cell line. Accordingly, this gene was designated as *MUM1* (multiple myeloma oncogene 1). The same probe was used to study the expression of the *MUM1* gene in various hematopoietic cell lines. The 6 kb message was expressed at high levels in most B cell lines and at low  
20     levels in peripheral T cell lines (Fig 4A). Cell lines derived from immature T cells, the myelomonocytic lineage, and erythroid lineage do not seem to express *MUM1*. In B cells, *MUM1* appears to be expressed throughout the development from the preB cell stage to the plasma cell stage (Fig 4B). However, some of the Burkitt's lymphoma derived cell lines such as BJA-B did not express this gene (data not shown). The expression level of the *MUM1* transcript in myeloma cell lines was also examined (Fig 4C).  
25     The SK-MM-1 cell line showed a 7.5-fold overexpression when compared with the other three IL-6 independent cell lines, suggesting a deregulated expression of the translocated allele. It is of interest that the IL-6 dependent XG-4, XG-7, and XG-10 cell lines are also expressing at high  
30     levels. Particularly, expression in the XG-7 cell line is 19.9 times the average of the aforementioned control cell lines.

### **MUMI cDNA cloning, sequencing, and homology search.**

Human spleen cDNA library was initially screened with a 2.1H probe followed by three times walking to 5' side using cDNA probes. A 5.5kb cDNA, approximately corresponding to the size detected by Northern analysis was isolated. This cDNA contained a 1,353 base pair open reading frame (ORF) and a long 3' untranslated region (Fig 5A). The ORF encodes for a protein of 451 amino acids with a predicted molecular weight of 50 kD (Fig 5B). The putative ATG initiation codon at position 217 has G at the -3 position which corresponds to the Kozak consensus sequence (23). The ORF is preceded by two in-frame stop codons. A database search demonstrated a significant similarity between MUM-1 ORF and the interferon regulatory factor (IRF) family proteins. The NH<sub>2</sub>-terminal of the MUM-1 ORF shares a high homology with all of the IRF family proteins which share a characteristic DNA binding motif consisting of the conserved 5 tryptophan residues (Fig 6A). The COOH-terminal also has a high homology with ICSBP (interferon consensus sequence binding protein)(21), ISGF3 $\gamma$  (interferon-stimulated gene factor-3 gamma)(22), and IRF-3 protein (23)(Fig 6B), although it did not have any homologous regions with IRF-1 and IRF-2 protein. The highest similarity (95.1%) and identity (91.8%) were found with a possible mouse homolog, LSIRF (lymphoid specific interferon regulatory factor)/Pip (PU-1 cofactor protein-1)(24,25). A high similarity was found with ICSBP (63.98%), ISGF3 $\gamma$  (55.8%), and IRF3 (50.1%) among the human IRF family protein members. A gene sequence encoding a nearly identical protein was recently deposited in GenBank. This gene, termed ICSAT (interferon consensus sequence binding protein in adult T-cell leukemia cell lines or activated T cells) is likely to be the same gene as MUM1



(26).

**Breakpoints at MUM1 locus in multiple myeloma.**

In order to analyze the exact location of the SK-MM-1  
5 breakpoint at the 6p25 locus and to explore the frequency of  
the MUM1 gene involvement in myeloma cases, we walked nearly  
55kb in a human placental genomic phage library around the  
MUM1 gene and determined the rough exon-intron structure as  
shown in Figure 7 (Fig. 7). The SK-MM-1 breakpoint was  
10 located 3' to the last exon, containing a poly A additional  
signal, consistent with an unaltered size of the MUM1  
transcript of this cell line in Northern analysis. Seven  
repeat-free genomic probes shown in Figure 7 have been used  
to investigate the rearrangement in Southern analyses of the  
15 11 MM cell lines and 18 MM cases. One case (case 10)  
displayed rearranged bands in *Bam*HI and *Xba*I digests when  
analyzed using a 0.9A probe located at 3' to the MUM1 gene.

**Cloning of the MUM2 locus from the U-266 multiple myeloma  
20 cell line.**

Using an experimental strategy analogous to the one  
described for the cloning of the MUM1 gene from the SK-MM-1  
cell line, a second genetic locus altered in multiple  
myeloma (MUM2) was identified by analyzing the U-266  
25 multiple myeloma cell line. Briefly, Southern blot analysis  
using *Bam*H restriction digestion and various Ig probes showed  
that U-266 DNA contained two rearranged fragments (shown by  
arrowheads in Fig. 9) containing C $\alpha$  sequences and lacking J  
sequences. These two fragments (der 14 and 14q32 in Fig. 9)  
30 were cloned from a genomic library constructed from U-266  
DNA along with a normal 14q 32 locus (14q32 germline in Fig.  
9). In order to determine whether a gene was located in

proximity to the chromosomal breakpoints in der 14, the 2.5 BE restriction fragment (see Fig. 9), which was at the opposite side of the Ig Ca sequences, was used to probe a Northern blot carrying RNA from various MM cell lines. The results (Fig. 10) showed that a 1.9 kb mRNA was detectable in some of these cell lines including U-266. This result showed that a gene, called MUM2, normally not present within the Ig locus on chromosome 14q32, had been translocated in proximity of the Ig locus in U-266 cells. Since the Ig locus contains strong transcriptional regulatory elements, it is likely that the expression of this gene is deregulated in these cells. The structure of the MUM2 gene and its protein are currently under investigation. The 2.5 BE probe and other probes derived from the der 14 phage can be used to screen MM cases for MUM2 rearrangements as shown for MUM1 (Fig. 7).

### **Experimental Discussion**

Using the experimental strategies used for the identification of the MUM1 and MUM2 genes in the SK-MM-1 and U-266 cell lines, respectively, it is possible to analyze most MM cases and isolate the corresponding genes. The scheme shown in Fig. 11 shows that the physiological IgH gene rearrangements (Fig. 11A) typically maintain linkage of C and J sequences and this linkage becomes detectable by using an appropriate restriction enzyme digestion (BamHI in the example in Fig. 11). Conversely, chromosomal translocations (14q+) affecting the IgH locus on 14q32 lead to breakage of the C-J linkage and the two sets of sequences appear on distinct restriction fragments. (Fig. 11B) Table 1 shows the application of this analysis to a panel of MM

cell lines and biopsies. The results show that at least 65% of cases show breakage of the C-J linkage within Ig J or switch regions. The restriction fragments containing either C or J sequences (R in Table 1) can be cloned as shown for the SK-MM-1 and U-266 cell lines and the genes flanking the chromosomal breakpoints can be used as probes to screen additional MM cases for similar rearrangements, whereas the sequence of the genes can be used to understand the consequences of these genetic lesions in multiple myeloma. Cloning of the chromosomal breakpoints and corresponding genes is currently ongoing for all of the MM cases shown in Table 1.

The method of analysis of 14q+ chromosomal translocations and identification of the genes altered in multiple myeloma of this invention will allow 1) the determination of chromosomal sequences involved in 14q+ translocations, the most important cytogenetic lesion associated with MM pathogenesis elucidation; 2) elucidation of specific gene lesions for MM; 3) a diagnostic method based on gene/DNA lesion and 4) a therapeutic approach aimed at counteracting the action of abnormal gene products.

Table 1. Summary of JH-C breakage analysis in MM cell lines and biopsies (cases). Rearrangement (R) involving physiologic Ig recombinations, i.e. retaining JH-C linkage are marked as R\*; rearrangements lacking JH-C linkage, and therefore suggesting a 14q+ chromosomal breakpoint, are marked as R. The latter represents candidates for cloning an further analysis.

Cell Line/Case	sIg	JH	Cμ	Cα	Sγ3'	possible breakpoint locus
10 RPMI-8226	λ	D/D	D/D	G	<u>R</u> /G	Sγ
U-266	Eλ	R/D	D/D	<u>R</u> / <u>R</u> /G	G	Sα
EJM	Gλ	<u>R</u> /R*	D/D	G	R*/G	JH~Sμ
XG-1	Aκ	R*/D	D/D	R*	G	ND
XG-2	Gλ	R*/D	D/D	G	R*/G	ND
15 XG-4	Gκ	R*/ <u>R</u>	D/D	G	R*/G	JH~Sμ
XG-5	λ	<u>R</u> /D	D/D	G	G	JH~Sμ
XG-6	Gλ	R*/ <u>R</u>	D/D	G	R*/G	JH~Sμ
XG-7	Aκ	R/D	D/D	R/G	<u>R</u> /D	Sγ
XG-10	G	R*/ <u>R</u>	D/D	G	R*/G	JH~Sμ
20 SK-MM-1	κ	R*/ <u>R</u>	<u>R</u> /D	G	R*/G	JH~Sμ
CASE125		R*	G	G	R*	ND
CASE33		<u>R</u> /R*	G	G	<u>R</u> /R*	Sγ
CASE34		R*	G	R*	G	ND
25 CASE93		R*	G	R*	G	ND
CASE91		R*	R*	<u>R</u>	G	Sα
CASE128		R*	G	R*	G	ND

R\*, comigrated bands with JH; R, target bands to isolate; ND, not determined

**Possible breakage in switch regions:**

Cell Lines	4/11 (36%)	
Cases	2/6 (33%)	Total 6/17 (35%)

**Possible breakage in JH ~ switch regions:**

Cell Lines	9/11 (82%)	
Cases	2/6 (33%)	Total 11/17 (65%)

**References**

1. Rabbitts, T.H. Chromosomal translocations in human cancer. Nature 372:143-149, 1994.  
5
2. Dalla-Favera, R., Bregni, M., Erickson, D., Patterson, D., Gallo, R.C., Croce, C.M. Human c-myc oncogene is located on the region of chromosome 8 that is translocated in Burkitt lymphoma cells. Proc Nat Acad Sci USA 79:7824, 1982.  
10
3. Tsujimoto, Y., Jaffe, E., Cossman, J., Gorham, J., Nowell, P.C., Croce, C.M. Clustering of breakpoints on chromosome 11 in human B-cell neoplasms with the t(11;14) chromosome translocation. Nature 315:340-343, 1985.  
15
4. Motokura, T., Bloom, T., Kim, H.G., Juppner, H., Ruderman, J.V., Kronenberg, H.M., Arnold, A. A novel cyclin encoded by a bcl-1 linked candidate oncogene. Nature 350:512-515, 1991.  
20
5. Tsujimoto, Y., Yunis, J., Onorato-Showe, L., Erikson, J., Nowell, P.C., Croce, C.M. Molecular cloning of chromosomal breakpoint of B-cell lymphomas and leukemias with the t(11;14) chromosome translocation. Science 224:1403-1406, 1984.  
25
6. Cleary, M.L., Sklar, J. : Nucleotide sequence of a t(14;18) chromosomal breakpoint in follicular lymphoma and demonstration of a breakpoint-cluster region near a transcriptionally active locus on chromosome 18. Proc  
30

Natl Acad Sci USA 82: 7439, 1985.

7. Bakhshi, A., Jensen, J.P., Goldman, P., Wright, J.J.,  
5 McBride, O.W., Epstein, A.L., Korsmeyer, S.J. Cloning  
the chromosomal breakpoint of t(14;18) human lymphomas:  
clustering around JH on chromosome 14 and near a  
transcriptional unit on 18. Cell 41:889, 1985.
8. Ye, B.H., Lista, F., Lo Coco, F., Knowles, D.M., Offit,  
10 K., Chaganti, R.S.K., Dalla-Favera, R. Alterations of  
a zinc finger-encoding gene, BCL-6, in diffuse  
large-cell lymphoma. Science 262:747-750, 1993.
9. Dewald, G.W., Kyle, R.A., Hicks, G.A., Greipp, P.R. The  
15 clinical significance of cytogenetic studies in 100  
patients with multiple myeloma, plasma cell leukemia,  
or amyloidosis. Blood 66:380-390, 1985.
10. Gould, J., Alexanian, R., Goodacre, A., Pathak, S.,  
20 Hecht, B., Barlogie, B. Plasma cell karyotype in  
multiple myeloma. Blood 71:453-456, 1988.
11. Weh, H.J., Gutensohn, K., Selbach, J., Kruse, R.,  
25 Wacker-Backhaus, G., Seeger, D., Fiedler, W., Fett, W.,  
Hossfeld, D.K. Karyotype in multiple myeloma and plasma  
cell leukemia. Eur J Cancer 29A:1269-1273, 1993.
12. Jernberg, H., Zech, L., Nilsson, K. Cytogenetic studies  
30 on human myeloma cell lines. Int J Cancer 40: 811-817,  
1987.
13. Zhang, X-G., Gaillard, J.P., Robillard, N., Lu, Z-Y.,

Gu, Z-G., Jourdan M., Boiron, J.M., Bataille, R., Klein, B. Reproducible obtaining of human myeloma cell lines as a model for tumor stem cell study in human multiple myeloma. Blood 83:3654-3663, 1994.

5

14. Seto, M., Yamamoto, K., Iida, S., Akao, Y., Utsumi, K.R., Kubonishi, I., Miyoshi, I., Ohtsuki, T., Yawata, Y., Namba, M., Motokura, T., Arnold, A., Takahashi, T., Ueda, R. Gene rearrangement and overexpression of PRAD1 in lymphoid malignancy with t(11;14)(q13;q32) translocation. Oncogene 7:1401 -1406, 1992.

10

15. Rabbitts, P.H., Douglas, J., Fisher, P., Nacheva, E., Karpas, A., Catovsky, D., Melo, J.V., Baer, R., Stinson, M.A., Rabbitts, T.H. Chromosome abnormalities at 11q13 in B cell tumours. Oncogene 3:99103, 1988.

15

16. Fiedler, W., Weh H.J., Hossfeld, D.K. Comparison of chromosome analysis and BCL-1 rearrangement in a series of patients with multiple myeloma. Br J Haematol 81: 58-61, 1992.

20

17. Taniwaki, M., Nishida, K., Takashima, T., Nakagawa, H., Fujii, H., Tamaki, T., Shimazaki, C., Horiike, S., Misawa, S., Kashima, K. Nonrandom chromosomal rearrangements of 14q32.3 and 19p13.3 and preferential deletion of lp in 21 patients with multiple myeloma and plasma cell leukemia. Blood 84: 2283-2290, 1994.

25

18. Sun, Z., Kitchingman, G.R. Sequencing of selected regions of the human immunoglobulin heavy-chain gene locus that completes the sequence from JH through delta

30

constant region. DNA sequence 1:347-355, 1991.

19. Eton, O., Scheinberg, D.A., Houghton, A.N.  
Establishment and characterization of two human myeloma  
cell lines secreting kappa light chains. *Leukemia* 3:  
729-735, 1989.
20. Mazars, G-R., Portier, M., Zhang, X-G., Jourdan, M.,  
Bataille, R., Theillet, C., Klein, B. Mutations of the  
pS3 gene in human myeloma cell lines. *Oncogene* 7:  
1015-1018, 1992.
21. Iida, S., Seto M., Yamamoto, K., Tojo, A., Asano, S.,  
Kamada, N., Ariyoshi, Y., Takahashi, T., Ueda, R. MLLT3  
gene on 9p22 involved in t(9;11) leukemia encodes a  
serine/proline rich protein homologous to MLLT1 on  
19p13. *Oncogene* 8(11):3085-3095, 1993.
22. Rao, P.H., Murty, V.V.V.S., Gaidano, G., Hauptschein,  
R., Dalla-Favera, R., and Chaganti, R.S.K. Subregional  
mapping of 8 single copy loci to chromosome 6 by  
fluorescence in situ hybridization. *Cytogenet. Cell  
Genet.* 66:272-273, 1994.
23. Kozak, M. The scanning model for translation: an  
update. *J. Cell. Biol.* 108:229-241, 1989.
24. Driggers, P.H., Ennist, D.L., Gleason, S.L., Mak, W-H.,  
Marka, M.S., Levi, B-Z., Flanagan, J.R., Appella, E.,  
Ozato, K. : *Proc Natl Acad Sci USA* 87: 3743-3747, 1990.
25. Veals, S.A., Schindler, C., Leonardo, D., Fu, X-Y.,



- 5       Aebersold, R., Damell, J.E., Levy, D.E. Subunit of an  
alpha-interferon-responsive transcription factor is  
related to interferon regulatory factor and myb  
families of DNA-binding proteins. Mol Cell Biol 12:  
3315-3324, 1992.
- 10       26. Grant, C.E., Vasa, M.Z., Deeley, R.G. cIRF-3, a new  
member of the interferon regulatory factor(IRF) family  
that is rapidly and transiently induced by dsRNA.  
Nucleic Acid Res 23:2137-2145, 1995.
- 15       27. Matsuyama, T., Grossman, A., Mittrucker, H-W.,  
Siderovski, D.P., Kiefer, F., Kawakami, T., Richardson,  
C.D., Taniguchi, T., Yoshinaga, S.K., Mak, T.W.  
Molecular cloning of LSIRF, a lymphoid-specific member  
of the interferon regulatory factor family that binds  
the interferon-stimulated response element(ISRE).  
Nucleic Acid Res 23:2127-2136, 1995.
- 20       28. Eisenbeis, C.F., Singh, H., Storb, U. Pip, a novel IRF  
family member, is a lymphoid-specific, PU.1-dependent  
transcriptional activator. Genes & Dev 9:1377-1387,  
1995.
- 25       29. Yamagata, T., Nishida, J., Tanaka, T., Sakai, R.,  
Mitani, K., Yoshida, M., Taniguchi, T., Yazaki, Y.,  
Hirai, H. A novel interferon regulatory factor family  
transcription factor, ICSAT/Pip/LSIRF, that negatively  
regulates the activity of interferon-regulated genes.  
30       Mol Cell Biol 16:1283-1294, 1996.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Dalla-Favera, Riccardo

(ii) TITLE OF INVENTION: IDENTIFICATION OF GENES ALTERED IN  
MULTIPLE MYELOMA

10

(iii) NUMBER OF SEQUENCES: 17

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: Cooper & Dunham LLP

(B) STREET: 1185 Avenue of the Americas

(C) CITY: New York

(D) STATE: New York

(E) COUNTRY: U.S.A.

(F) ZIP: 10036

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

25

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE: 28-MAY-1996

30

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: White, John P.

(B) REGISTRATION NUMBER: 28,678

35

(C) REFERENCE/DOCKET NUMBER: 50995

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (212) 278-0400

(B) TELEFAX: (212) 391-0525

40

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile Asp Ser Gly Lys Tyr Pro  
1 5 10 15  
Gly Leu Val Trp Glu Asn Glu Glu Lys Ser Ile Phe Arg Ile Pro Trp  
20 25 30  
Lys His Ala Gly Lys Gln Asp Tyr Asn Arg Glu Glu Asp Ala Ala Leu  
35 40 45  
Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys Phe Arg Glu Gly Ile Asp  
50 55 60  
Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg Leu Arg Cys Ala Leu Asn  
65 70 75 80  
Lys Ser Asn Asp Phe Glu Glu Leu Val Glu Arg Ser Gln Leu Asp Ile  
85 90 95  
Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val Pro Glu  
100 105

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

10 Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile Asp Ser Gly Lys Tyr Pro  
1 5 10 15

Gly Leu Val Trp Glu Asn Glu Glu Lys Ser Val Phe Arg Ile Pro Trp  
20 25 30

15 Lys His Ala Gly Lys Gln Asp Tyr Asn Arg Glu Glu Asp Ala Ala Leu  
35 40 45

Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys Phe Arg Glu Gly Ile Asp  
20 50 55 60

Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg Leu Arg Cys Ala Leu Asn  
65 70 75 80

25 Lys Ser Asn Asp Phe Glu Glu Leu Val Glu Arg Ser Gln Leu Asp Ile  
85 90 95

Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val Pro Glu  
100 105

30

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5 Arg Met Arg Pro Trp Leu Glu Met Gln Ile Asn Ser Asn Gln Ile Pro  
1 5 10 15

Gly Leu Ile Trp Ile Asn Lys Glu Glu Met Ile Phe Gln Ile Pro Trp  
20 25 30

10 Lys His Ala Ala Lys His Gly Trp Asp Ile Asn Lys Asp Ala Cys Leu  
35 40 45

Phe Arg Ser Trp Ala Ile His Thr Gly Arg Tyr Lys Ala Gly Glu Lys  
50 55 60

15 Glu Pro Asp Pro Lys Thr Trp Lys Ala Asn Phe Arg Cys Ala Met Asn  
65 70 75 80

Ser Leu Pro Asp Ile Glu Glu Val Lys Asp Gln Lys Arg Asn Lys Gly  
85 90 95

20 Ser Ser Ala Val Arg Val Tyr Arg Met Leu Pro Pro  
100 105

(2) INFORMATION FOR SEQ ID NO:4:

25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 108 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

40 Arg Met Arg Pro Trp Leu Glu Glu Gln Ile Asn Ser Asn Thr Ile Pro  
1 5 10 15

Gly Leu Lys Trp Leu Asn Lys Glu Lys Lys Ile Phe Gln Ile Pro Trp

20 25 30

Met His Ala Ala Arg His Gly Trp Asp Val Glu Lys Asp Ala Pro Leu  
35 40 45

5

Phe Arg Asn Trp Ala Ile His Thr Gly Lys His Gln Pro Gly Val Asp  
50 55 60

Lys Pro Asp Pro Lys Thr Trp Lys Ala Asn Phe Arg Cys Ala Met Asn  
10 65 70 75 80

Ser Leu Pro Asp Ile Glu Glu Val Lys Asp Lys Ser Ile Lys Lys Gly  
85 90 95

15

Asn Asn Ala Phe Arg Val Tyr Arg Met Leu Pro Leu  
100 105

(2) INFORMATION FOR SEQ ID NO:5:

20 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 107 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Leu Arg Gln Trp Leu Ile Glu Gln Ile Asp Ser Ser Met Tyr Pro  
1 5 10 15

35

Gly Leu Ile Trp Glu Asn Glu Glu Lys Ser Met Phe Arg Ile Pro Trp  
20 25 30

Lys His Ala Gly Lys Gln Asp Tyr Asn Gln Glu Val Asp Ala Ser Ile  
40 35 40 45

Phe Lys Ala Trp Ala Val Phe Lys Gly Lys Phe Lys Glu Gly Asp Lys

	50		55		60											
	Ala	Glu	Pro	Ala	Thr	Trp	Lys	Thr	Arg	Leu	Arg	Cys	Ala	Leu	Asn	Lys
	65					70					75					80
5	Ser	Pro	Asp	Phe	Glu	Glu	Val	Thr	Asp	Arg	Ser	Gln	Leu	Asp	Ile	Ser
				85						90					95	
	Glu	Pro	Tyr	Lys	Val	Tyr	Arg	Ile	Val	Pro	Glu					
10				100					105							

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 107 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: peptide

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

	Lys	Leu	Arg	Asn	Trp	Val	Val	Glu	Gln	Val	Glu	Ser	Gly	Gln	Phe	Pro
	1				5					10					15	
30	Gly	Val	Cys	Trp	Asp	Asp	Thr	Ala	Lys	Thr	Met	Phe	Arg	Ile	Pro	Trp
				20					25					30		
	Lys	His	Ala	Gly	Lys	Gln	Asp	Phe	Arg	Glu	Asp	Gln	Asp	Ala	Ala	Phe
			35					40					45			
35	Phe	Lys	Ala	Trp	Ala	Ile	Phe	Lys	Gly	Lys	Tyr	Lys	Glu	Gly	Asp	Thr
		50					55					60				
	Gly	Gly	Pro	Ala	Val	Trp	Lys	Thr	Arg	Leu	Arg	Cys	Ala	Leu	Asn	Lys
40	65					70					75					80
	Ser	Ser	Glu	Phe	Lys	Glu	Val	Pro	Glu	Arg	Gly	Arg	Met	Asp	Val	Ala

85

90

95

Glu Pro Tyr Lys Val Tyr Gln Leu Leu Pro Pro  
100 105

5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 107 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

20

Arg Ile Leu Pro Trp Leu Val Ser Gln Leu Asp Leu Gly Gln Leu Glu  
1 5 10 15

25

Gly Val Ala Trp Val Asn Lys Ser Arg Thr Arg Phe Arg Ile Pro Trp  
20 25 30

Lys His Gly Leu Arg Gln Asp Ala Gln Gln Glu Asp Phe Gly Ile Phe  
35 40 45

30

Gln Ala Trp Ala Glu Ala Thr Gly Ala Tyr Val Pro Gly Arg Asp Lys  
50 55 60

35

Pro Asp Leu Pro Thr Trp Lys Arg Asn Phe Arg Ser Ser Ala Leu Asn  
65 70 75 80

Arg Lys Glu Gly Leu Arg Leu Ala Glu Asp Arg Ser Lys Asp Pro His  
85 90 95

40

Asp Pro His Lys Ile Tyr Glu Phe Val Asn Ser  
100 105

(2) INFORMATION FOR SEQ ID NO:8:



(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 95 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Lys	Arg	Leu	Cys	Gln	Ser	Thr	Ile	Tyr	Trp	Asp	Gly	Pro	Leu	Ala	Leu
1			5					10					15		
Cys	Asn	Asp	Arg	Pro	Asn	Lys	Leu	Glu	Arg	Asp	Gln	Thr	Cys	Lys	Leu
			20					25					30		
Phe	Asp	Thr	Gln	Gln	Phe	Leu	Ser	Glu	Leu	Gln	Ala	Phe	Ala	His	His
			35				40					45			
Gly	Arg	Ser	Leu	Pro	Arg	Phe	Gln	Val	Thr	Leu	Cys	Phe	Gly	Glu	Glu
			50				55				60				
Phe	Pro	Asp	Pro	Gln	Arg	Gln	Arg	Lys	Leu	Ile	Thr	Ala	His	Val	Glu
65					70				75						80
Pro	Leu	Leu	Ala	Arg	Gln	Leu	Tyr	Tyr	Phe	Ala	Gln	Gln	Asn	Ser	
					85				90					95	

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 95 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5	Lys	Arg	Leu	Cys	Gln	Ser	Arg	Ile	Tyr	Trp	Asp	Gly	Pro	Leu	Ala	Leu
	1				5					10					15	
	Cys	Ser	Asp	Arg	Pro	Asn	Lys	Leu	Glu	Arg	Asp	Gln	Thr	Cys	Lys	Leu
				20					25					30		
10	Phe	Asp	Thr	Gln	Gln	Phe	Leu	Ser	Glu	Leu	Gln	Val	Phe	Ala	His	His
			35					40					45			
	Gly	Arg	Pro	Ala	Pro	Arg	Phe	Gln	Val	Thr	Leu	Cys	Phe	Gly	Glu	Glu
		50					55					60				
15	Phe	Pro	Asp	Pro	Gln	Arg	Gln	Arg	Lys	Leu	Ile	Thr	Ala	His	Val	Glu
	65					70					75					80
	Pro	Leu	Leu	Ala	Arg	Gln	Leu	Tyr	Tyr	Phe	Ala	Gln	Gln	Asn	Thr	
20					85					90					95	

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 96 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: peptide

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

	Lys	Arg	Leu	Cys	Gln	Gly	Arg	Val	Phe	Cys	Ser	Gly	Asn	Ala	Val	Val
	1				5					10					15	
40	Cys	Lys	Gly	Arg	Pro	Asn	Lys	Leu	Glu	Arg	Asp	Glu	Val	Val	Gln	Val
				20					25					30		

Phe Asp Thr Ser Gln Phe Phe Arg Glu Leu Gln Gln Phe Tyr Asn Ser  
35 40 45

5 Gln Gly Arg Leu Pro Asp Gly Arg Val Val Leu Cys Phe Gly Glu Glu  
50 55 60

Phe Pro Asp Met Ala Pro Leu Arg Ser Lys Leu Ile Leu Val Gln Ile  
65 70 75 80

10 Glu Gln Leu Tyr Val Arg Gln Leu Ala Glu Glu Ala Gly Lys Ser Cys  
85 90 95

(2) INFORMATION FOR SEQ ID NO:11:

15

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 96 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

20

- (ii) MOLECULE TYPE: peptide

25

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

30

Gln Arg Leu Cys Pro Ile Pro Ile Ser Trp Asn Ala Pro Gln Ala Pro  
1 5 10 15

Pro Gly Pro Gly Pro His Leu Leu Pro Ser Asn Glu Cys Val Glu Leu  
20 25 30

35

Phe Arg Thr Ala Tyr Phe Cys Arg Asp Leu Val Arg Tyr Phe Gln Gly  
35 40 45

Leu Gly Pro Pro Pro Lys Phe Gln Val Thr Leu Asn Phe Trp Glu Glu  
50 55 60

40

Ser His Gly Ser Ser His Thr Pro Gln Asn Leu Ile Thr Val Lys Met  
65 70 75 80

Glu Gln Ala Phe Ala Arg Tyr Leu Leu Glu Gln Thr Pro Glu Gln Gln  
85 90 95

5 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 100 amino acids

(B) TYPE: amino acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

20 Gln Arg Leu Gly His Cys His Thr Tyr Trp Ala Val Ser Glu Glu Leu  
1 5 10 15

Leu Pro Asn Ser Gly His Gly Pro Asp Gly Glu Val Pro Lys Asp Lys  
20 25 30

25

Glu Gly Gly Val Phe Asp Leu Gly Pro Phe Ile Val Asp Leu Ile Thr  
35 40 45

30

Phe Thr Glu Gly Ser Gly Arg Ser Pro Arg Tyr Ala Trp Leu Phe Cys  
50 55 60

Val Gly Glu Ser Trp Pro Gln Asp Gln Pro Trp Thr Lys Arg Leu Val  
65 70 75 80

35

Met Val Lys Val Val Pro Thr Cys Leu Arg Ala Leu Val Glu Met Ala  
85 90 95

Arg Val Gly Gly  
100

40

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5176 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: other nucleic acid

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 217..1569

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

15

GCCTGACCAA CATGGTAAAA CCCCATCTCT GCTAAACTA CAAAAATTA GCTGGATGTG  
60

20

GTGGCAGGGA ACCTGTCATC CCAGCTAGTT GGGAGACTGA GGCAGGAGAA TCGCTCGATC  
120

TTGGGACCCA CCGCTGCCCT CAGCTCCGAG TCCAGGGCGA GTGCAGAGCA CAGCGGGCGG  
180

25

AGGACCCCGG GCGCGGGCGC GGACGGCACG CGGGGC ATG AAC CTG GAG GGC GGC  
234

Met Asn Leu Glu Gly Gly  
1 5

30

GGC CGA GGC GGA GAG TTC GGC ATG AGC GCG GTG AGC TGC GGC AAC GGC  
282

Gly Arg Gly Gly Glu Phe Gly Met Ser Ala Val Ser Cys Gly Asn Gly  
10 15 20

35

AAG CTC CGC CAG TGG CTG ATC GAC CAG ATC GAC AGC GGC AAG TAC CCC  
330

Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile Asp Ser Gly Lys Tyr Pro  
25 30 35

40

GGG CTG GTG TGG GAG AAC GAG GAG AAG AGC ATC TTC CGC ATC CCC TGG  
378

Gly Leu Val Trp Glu Asn Glu Glu Lys Ser Ile Phe Arg Ile Pro Trp  
40 45 50

5 AAG CAC GCG GGC AAG CAG GAC TAC AAC CGC GAG GAG GAC GCC GCG CTC  
426

Lys His Ala Gly Lys Gln Asp Tyr Asn Arg Glu Glu Asp Ala Ala Leu  
55 60 65 70

10 TTC AAG GCT TGG GCA CTG TTT AAA GGA AAG TTC CGA GAA GGC ATC GAC  
474

Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys Phe Arg Glu Gly Ile Asp  
75 80 85

15 AAG CCG GAC CCT CCC ACC TGG AAG ACG CGC CTG CGG TGC GCT TTG AAC  
522

Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg Leu Arg Cys Ala Leu Asn  
90 95 100

20 AAG AGC AAT GAC TTT GAG GAA CTG GTT GAG CGG AGC CAG CTG GAC ATC  
570

Lys Ser Asn Asp Phe Glu Glu Leu Val Glu Arg Ser Gln Leu Asp Ile  
105 110 115

25 TCA GAC CCG TAC AAA GTG TAC AGG ATT GTT CCT GAG GGA GCC AAA AAA  
618

Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val Pro Glu Gly Ala Lys Lys  
120 125 130

30 GGA GCC AAG CAG CTC ACC CTG GAG GAC CCG CAG ATG TCC ATG AGC CAC  
666

Gly Ala Lys Gln Leu Thr Leu Glu Asp Pro Gln Met Ser Met Ser His  
135 140 145 150

35 CCC TAC ACC ATG ACA ACG CCT TAC CCT TCG CTC CCA GCC CAG CAG GTT  
714

Pro Tyr Thr Met Thr Thr Pro Tyr Pro Ser Leu Pro Ala Gln Gln Val  
155 160 165

40 CAC AAC TAC ATG ATG CCA CCC CTC GAC CGA AGC TGG AGG GAC TAC GTC  
762

His Asn Tyr Met Met Pro Pro Leu Asp Arg Ser Trp Arg Asp Tyr Val  
170 175 180

CCG GAT CAG CCA CAC CCG GAA ATC CCG TAC CAA TGT CCC ATG ACG TTT  
810

Pro Asp Gln Pro His Pro Glu Ile Pro Tyr Gln Cys Pro Met Thr Phe  
185 190 195

5

GGA CCC CGC GGC CAC CAC TGG CAA GGC CCA GCT TGT GAA AAT GGT TGC  
858

Gly Pro Arg Gly His His Trp Gln Gly Pro Ala Cys Glu Asn Gly Cys  
200 205 210

10

CAG GTG ACA GGA ACC TTT TAT GCT TGT GCC CCA CCT GAG TCC CAG GCT  
906

Gln Val Thr Gly Thr Phe Tyr Ala Cys Ala Pro Pro Glu Ser Gln Ala  
215 220 225 230

15

CCC GGA GTC CCC ACA GAG CCA AGC ATA AGG TCT GCC GAA GCC TTG GCG  
954

Pro Gly Val Pro Thr Glu Pro Ser Ile Arg Ser Ala Glu Ala Leu Ala  
235 240 245

20

TTC TCA GAC TGC CGG CTG CAC ATC TGC CTG TAC TAC CGG GAA ATC CTC  
1002

Phe Ser Asp Cys Arg Leu His Ile Cys Leu Tyr Tyr Arg Glu Ile Leu  
250 255 260

25

GTG AAG GAG CTG ACC ACG TCC AGC CCC GAG GGC TGC CGG ATC TCC CAT  
1050

Val Lys Glu Leu Thr Thr Ser Ser Pro Glu Gly Cys Arg Ile Ser His  
265 270 275

30

GGA CAT ACG TAT GAC GCC AGC AAC CTG GAC CAG GTC CTG TTC CCC TAC  
1098

Gly His Thr Tyr Asp Ala Ser Asn Leu Asp Gln Val Leu Phe Pro Tyr  
280 285 290

35

CCA GAG GAC AAT GGC CAC AGG AAA AAC ATT GAG AAC CTG CTG AGC CAC  
1146

Pro Glu Asp Asn Gly His Arg Lys Asn Ile Glu Asn Leu Leu Ser His  
295 300 305 310

40

CTG GAG AGG GGC GTG GTC CTC TGG ATG GCC CCC GAC GGG CTC TAT GCG  
1194

Leu Glu Arg Gly Val Val Leu Trp Met Ala Pro Asp Gly Leu Tyr Ala  
315 320 325

5 AAA AGA CTG TGC CAG AGC ACG ATC TAC TGG GAC GGG CCC CTG GCG CTG  
1242

Lys Arg Leu Cys Gln Ser Thr Ile Tyr Trp Asp Gly Pro Leu Ala Leu  
330 335 340

10 TGC AAC GAC CGG CCC AAC AAA CTG GAG AGA GAC CAG ACC TGC AAG CTC  
1290

Cys Asn Asp Arg Pro Asn Lys Leu Glu Arg Asp Gln Thr Cys Lys Leu  
345 350 355

15 TTT GAC ACA CAG CAG TTC TTG TCA GAG CTG CAA GCG TTT GCT CAC CAC  
1338

Phe Asp Thr Gln Gln Phe Leu Ser Glu Leu Gln Ala Phe Ala His His  
360 365 370

20 GGC CGC TCC CTG CCA AGA TTC CAG GTG ACT CTA TGC TTT GGA GAG GAG  
1386

Gly Arg Ser Leu Pro Arg Phe Gln Val Thr Leu Cys Phe Gly Glu Glu  
375 380 385 390

25 TTT CCA GAC CCT CAG AGG CAA AGA AAG CTC ATC ACA GCT CAC GTA GAA  
1434

Phe Pro Asp Pro Gln Arg Gln Arg Lys Leu Ile Thr Ala His Val Glu  
395 400 405

30 CCT CTG CTA GCC AGA CAA CTA TAT TAT TTT GCT CAA CAA AAC AGT GGA  
1482

Pro Leu Leu Ala Arg Gln Leu Tyr Tyr Phe Ala Gln Gln Asn Ser Gly  
410 415 420

35 CAT TTC CTG AGG GGC TAC GAT TTA CCA GAA CAC ATC AGC AAT CCA GAA  
1530

His Phe Leu Arg Gly Tyr Asp Leu Pro Glu His Ile Ser Asn Pro Glu  
425 430 435

40 GAT TAC CAC AGA TCT ATC CGC CAT TCC TCT ATT CAA GAA TGAAAAATGT  
1579

Asp Tyr His Arg Ser Ile Arg His Ser Ser Ile Gln Glu  
440 445 450



CAAGATGAGT GGT TTTCTTTT TTCCTTTTTT TTTT TTTT TTTTGATACG GAGATACGGG  
1639

5 GTCTTGCTCT GTCTCCCAGG CTGGAGTGCA GTGACACAAT CTCAGCTCAC TGTGACCTCC  
1699

GCCTCCTGGG TTCAAGAGAC TCTCCTGCCT CAGCCTCCCT GGTAGCTGGG ATTACAGGTG  
1759

10 TGAGCCACTG CACCCACCCA AGACAAGTGA TTTTCATTGT AAATATTTGA CTTTAGTGAA  
1819

AGCGTCCAAT TGA CTGCCCT CTTACTGTTT TGAGGAACTC AGAAGTGGAG ATTTCA GTTC  
1879

15 AGCGGTTGAG GAGAATTGCG GCGAGACAAG CATGGAAAAT CAGTGACATC TGATTGGCAG  
1939

20 ATGAGCTTAT TTCAAAGGA AGGGTGGCTT TGCATTTTCT TGTGTTCTGT AGACTGCCAT  
1999

CATTGATGAT CACTGTGAAA ATTGACCAAG TGATGTGTTT ACATTTACTG AAATGCGCTC  
2059

25 TTAAATTTGT TGTAGATTAG GTCTTGCTGG AAGACAGAGA AAAGTGCCT TTCAGTATTG  
2119

ACACTGACTA GAGTGATGAC TGCTTG TAGG TATGTCTGTG CCATTTCTCA GGGAAGTAAG  
2179

30 ATGTAAATTG AAGAAGCCTC ACACGTAAAA GAAATGTATT AATGTATGTA GGAGCTGCAG  
2239

35 TTCTTGTGGA AGACACTTGC TGAGTGAAGG AAATGAATCT TTGACTGAAG CCGTGCCTGT  
2299

AGCCTTG GGG AGGCCCATCC CCCACCTGCC AGCGGTTTCC TGGTGTGGGT CCCTCTGCCC  
2359

40 CACCCTCCTT CCCATTGGCT TTCTCTCCTT GGCCTTTCCT GGAAGCCAGT TAGTAACTT  
2419

CCTATTTTCT TGAGTCAAAA AACATGAGCG CTACTCTTGG ATGGGACATT TTTGTCTGTC  
2479

5 CTACAATCTA GTAATGTCTA AGTAATGGTT AAGTTTTCTT GTTCTGTCAT CTTTTTGACC  
2539

CTCATTCCTT AGAGATGCTA AAATTCCTCG CATAAAGAAG AAGAAATTAA GGAACATAAA  
2599

10 TCTTAATACT TGAAGTGTG CCCTTCTGTC CAAGTACTTA ACTATCTGTT CCCTTCCTCT  
2659

GTGCCACGCT CCTCTGTTTG TTTGGCTGTC CAGCGATCAG CCATGGCGAC ACTAAAGGAG  
2719

15 GAGGAGCCGG GGAAGTCCAG GCTGGAGAGC ACTGCCAGGA CCCACCACTG GAAGCAGGAT  
2779

GGAGCTGACT ACGGAAGTGC AACTCAGTG GGCTGTTTCT GCTTATTTCA TCTGTTCTAT  
2839

20 GCTTCCTCGT GCCAATTATA GTTTGACAGG GCCTTAAAT TACTTGGCTT TTTCCAAATG  
2899

25 CTTCTATTTA TAGAAATCCC AAAGACCTCC ACTTGCTTAA GTATACCTAT CACTTACATT  
2959

TTTGTGGTTT TGAGAAAGTA CAGCAGTAGA CTGGGGCGTC ACCTCCAGGC CGTTTCTCAT  
3019

30 ACTACAGGAT ATTTACTATT ACTCCCAGGA TTCAGCAGAA GATTGCGTTA GCTCTCAAAT  
3079

GTGTGTTCCCT GCTTTTCTAA TGGATATTTT AAATTCATT CACAAGCACC TAGTAAGTGC  
3139

35 CTGCTGTATC CCTACATTAC ACAGTTCAGC CTTTATCAAG CTTAGTGAGC AGTGAGCACT  
3199

40 GAAACATTAT TTTTAAATGT TTAATAAGTT TCTAATATTA AAGTCAGAAT ATTAATACAA  
3259

TTAATATTAA TATTAACTAC AGAAAAGACA AACAGTAGAG AACAGCAAAA AAATAAAAAG  
3319

5 GATCTCCTTT TTTCCCAGCC CAAATTCTCC TCTCTAAAAG TGTCCACAAG AAGGGGTGTT  
3379

TATTCTTCCA ACACATTTCA CTTTTCTGTA AATATACATA AACTTAAAAA GAAAACCTCA  
3439

10 TGGAGTCATC TTGCACACAC TTTTCATGCA GTGCTCTTTG TAGCTAAACA GTGAAGATTT  
3499

15 ACCTCGTTCT GCTCAGAGGC CTTGCTGTGG AGCTCCACTG CCATGTACCC AGTAGGGTTT  
3559

GACATTTTCAT TAGCCATGCA ACATGGATAT GTATTGGGCA GCAGACTGTG TTTTCGTGAAC  
3619

20 TGCAGTGATG TATACATCTT ATAGATGCAA AGTATTTTGG GGTATATTAT CCTAAGGGAA  
3679

GATAAAGATG ATATTAAGAA CTGCTGTTTC ACGGGGCCCT TACCTGTGAC CCTCTTTGCT  
3739

25 GAAGAATATT TAACCCACACA CAGCACTTCA AAGAAGCTGT CTTGGAAGTC TGTCTCAGGA  
3799

GCACCCTGTC TTCTTAATTC TCCAAGCGGA TGCTCCATTT CAATTGCTTT GTGACTTCTT  
3859

30 CTTCTTTGTT TTTTAAATA TTATGCTGCT TTAACAGTGG AGCTGAATTT TCTGGAAAAT  
3919

35 GCTTCTTGGC TGGGGCCACT ACCTCCTTTC CTATCTTTAC ATCTATGTGT ATGTTGACTT  
3979

TTTAAAATTC TGAGTGATCC AGGGTATGAC CTAGGGAATG AACTAGCTAT GGAAATAACT  
4039

40 CAGGGTTAGG AATCCTAGCA CTTGTCTCAG GACTCTGAAA AGGAACGGCT TCCTCATGCC  
4099

TTGTCTTGAT AAAGTGGAAT TGGCAAATA GAATTTAGTT TGTACTCAGT GGACAGTGCT  
4159

5 GTTGAAGATT TGAGGACTTG TTAAAGAGCA CTGGGTCATA TGGAAAAAAT GTATGTGTCT  
4219

CCCCAGGTGC ATTTTCTTGG TTTATGTCTT GTTCTTGAGA TTTTGTATAT TTAGGAAAAC  
4279

10 CTCAAGCAGT AATTAATATC TCCTGGAACA CTATAGAGAA CCAAGTGACC GACTCATTTA  
4339

CAACTGAAAC CTAGGAAGCC CCTGAGTCCT GAGCGAAAAC AGGAGAGTTA GTCGCCCTAC  
4399

15 AGAAAACCCA GCTAGACTAT TGGGTATGAA CTAAAAAGAG ACTGTGCCAT GGTGAGAAAA  
4459

ATGTAAAATC CTACAGTGGA ATGAGCAGCC CTTACAGTGT TGTTACCACC AAGGGCAGGT  
4519

20 AGGTATTAGT GTTTGAAAAA GCTGGTCTTT GAGCGAGGGC ATAAATACAG CTAGCCCCAG  
4579

25 GGGTGGAAACA ACTGTGGGAG TCTTGGGTAC TCGCACCTCT TGGCTTTGTT GATGCTCCGC  
4639

CAGGAAGGCC ACTTGTGTGT GCGTGTCTAGT TACTTTTTTA GTAACAATTC AGATCCAGTG  
4699

30 TAAACTTCCG TTCATTGCTC TCCAGTCACA TGCCCCCACT TCCCCACAGG TGAAAGTTTT  
4759

TCTGAAGTGT TGGGATTGGT TAAGGTCTTT ATTTGTATTA CGTATCTCCC CAAGTCCTCT  
4819

35 GTGGCCAGCT GCATCTGTCT GAATGGTGCG TGAAGGCTCT CAGACCTTAC ACACCATTTT  
4879

40 GTAAGTTATG TTTTACATGC CCCGTTTTTG AGACTGATCT CGATGCAGGT GGATCTCCTT  
4939

GAGATCCTGA TAGCCTGTTA CAGGAATGAA GTAAAGGTCA GTTTTTTTTG TATTGATTTT  
4999

5 CACAGCTTTG AGGAACATGC ATAAGAAATG TAGCTGAAGT AGAGGGGACG TGAGAGAAGG  
5059

GCCAGGCCGG CAGGCCAACC CTCCTCCAAT GGAAATTCCC GTGTTGCTTC AACTGAGAC  
5119

10 AGATGGGACT TAACAGGCAA TGGGGTCCAC TTCCCCCTCT TCAGCATCCC CCGTACC  
5176

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 451 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

25 Met Asn Leu Glu Gly Gly Gly Arg Gly Gly Glu Phe Gly Met Ser Ala  
1 5 10 15  
Val Ser Cys Gly Asn Gly Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile  
20 25 30  
30 Asp Ser Gly Lys Tyr Pro Gly Leu Val Trp Glu Asn Glu Glu Lys Ser  
35 40 45  
Ile Phe Arg Ile Pro Trp Lys His Ala Gly Lys Gln Asp Tyr Asn Arg  
35 50 55 60  
Glu Glu Asp Ala Ala Leu Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys  
65 70 75 80  
40 Phe Arg Glu Gly Ile Asp Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg  
85 90 95

	Leu Arg Cys Ala Leu Asn Lys Ser Asn Asp Phe Glu Glu Leu Val Glu	
	100	105 110
5	Arg Ser Gln Leu Asp Ile Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val	
	115	120 125
	Pro Glu Gly Ala Lys Lys Gly Ala Lys Gln Leu Thr Leu Glu Asp Pro	
	130	135 140
10	Gln Met Ser Met Ser His Pro Tyr Thr Met Thr Thr Pro Tyr Pro Ser	
	145	150 155 160
	Leu Pro Ala Gln Gln Val His Asn Tyr Met Met Pro Pro Leu Asp Arg	
	165	170 175
15	Ser Trp Arg Asp Tyr Val Pro Asp Gln Pro His Pro Glu Ile Pro Tyr	
	180	185 190
	Gln Cys Pro Met Thr Phe Gly Pro Arg Gly His His Trp Gln Gly Pro	
20	195	200 205
	Ala Cys Glu Asn Gly Cys Gln Val Thr Gly Thr Phe Tyr Ala Cys Ala	
	210	215 220
25	Pro Pro Glu Ser Gln Ala Pro Gly Val Pro Thr Glu Pro Ser Ile Arg	
	225	230 235 240
	Ser Ala Glu Ala Leu Ala Phe Ser Asp Cys Arg Leu His Ile Cys Leu	
	245	250 255
30	Tyr Tyr Arg Glu Ile Leu Val Lys Glu Leu Thr Thr Ser Ser Pro Glu	
	260	265 270
	Gly Cys Arg Ile Ser His Gly His Thr Tyr Asp Ala Ser Asn Leu Asp	
35	275	280 285
	Gln Val Leu Phe Pro Tyr Pro Glu Asp Asn Gly His Arg Lys Asn Ile	
	290	295 300
40	Glu Asn Leu Leu Ser His Leu Glu Arg Gly Val Val Leu Trp Met Ala	
	305	310 315 320

Pro Asp Gly Leu Tyr Ala Lys Arg Leu Cys Gln Ser Thr Ile Tyr Trp  
325 330 335

5 Asp Gly Pro Leu Ala Leu Cys Asn Asp Arg Pro Asn Lys Leu Glu Arg  
340 345 350

Asp Gln Thr Cys Lys Leu Phe Asp Thr Gln Gln Phe Leu Ser Glu Leu  
355 360 365

10 Gln Ala Phe Ala His His Gly Arg Ser Leu Pro Arg Phe Gln Val Thr  
370 375 380

Leu Cys Phe Gly Glu Glu Phe Pro Asp Pro Gln Arg Gln Arg Lys Leu  
385 390 395 400

15 Ile Thr Ala His Val Glu Pro Leu Leu Ala Arg Gln Leu Tyr Tyr Phe  
405 410 415

Ala Gln Gln Asn Ser Gly His Phe Leu Arg Gly Tyr Asp Leu Pro Glu  
20 420 425 430

His Ile Ser Asn Pro Glu Asp Tyr His Arg Ser Ile Arg His Ser Ser  
435 440 445

25 Ile Gln Glu  
450

(2) INFORMATION FOR SEQ ID NO:15:

30 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 152 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TTTTCTCTAC AGTCACCTCC CTGTTTACCA AAGATAATCA CAATAAGTCC AGTTTACTTA  
60

5 CAAAACAAGT TTAGTTATTA GAGGAACTA AAACCTCAGG ATTCAGTCCA GATAATTTTT  
120

AAAACTCTA AAACAATGGA CAGGGCTAGA AT  
152

10 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 152 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

25 TGGGCTCGGC CTGGTGGGGC AGCCACAGCG GGACGCAGTA GTGAAAGTCC AGTTTACTTA  
60

CAAAAACAAGT TTAGTTATTA GAGGAACTA AAACCTCAGG ATTCAGCAGG GCATGAGGAG  
120

30

GCAGCTCCTC ACCCTCCCTT TCTCTTTTGT AC  
152

(2) INFORMATION FOR SEQ ID NO:17:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 152 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)



(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

5 TGGGCTCGGC CTTGGTGGGG CAGCCACAGC GGGACGCAAG TAGTGAGGGC ACTCAGAACG  
60  
CCACTCAGCC CCGACAGGGC ACTCAGAACG CCACTCAGCC CCGACAGGCA GGGCACGAGG  
10 120  
AGGCAGCTCC TCACCCTCCC TTTCTCTTTT GT  
152

15

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207

What is claimed is:

1. A method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of:
  - (a) obtaining a DNA sample from the subject suffering from multiple myeloma;
  - (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample;
  - (c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction;
  - (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene;
  - (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones from step (d);
  - (f) hybridizing said fluorescent probes with metaphase chromosomes; and
  - (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma.
2. The method of claim 1, wherein step (b) is performed by Southern blotting.

3. The method of claim 1, wherein step (b) is performed by polymerase chain reaction with appropriate probes.
4. The method of claim 1, wherein the genomic library is  
5 a phage vector library.
5. The method of claim 4, wherein the genomic DNA fragments are generated by cleaving genomic DNA from cells of the subject with an appropriate restriction  
10 enzyme.
6. The method of claim 5, wherein the restriction enzyme is *Bam*HI.
7. The method of claim 5, wherein the restriction enzyme  
15 is *Sau*3AI.
8. The method of claim 1, wherein the probe of step (d) is a human IgH J region JH probe.
9. The method of claim 1, wherein the probe of step (d) is  
20 a human IgH C $\mu$  probe.
10. The method of claim 1, wherein the probe of step (d) is  
25 a human IgH C $\gamma$ 2 probe.
11. The method of claim 1, wherein the chromosomal breakpoint identified is a t(6;14)(p25;q32) translocation.
12. The method of claim 1, wherein the chromosomal  
30 breakpoint identified is a t(14;15) translocation.

13. A method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma which comprises steps of:

- a) selecting a probe having a sequence of a chromosome other than chromosome 14, identified at the chromosomal breakpoint detected in a subject suffering from multiple myeloma, wherein said probe is capable of hybridizing to the unique sequence of the gene other than the immunoglobulin gene altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma;
- b) contacting said probe with mRNA isolated from a cell under conditions permitting formation of a complex between said probe and the mRNA;
- c) isolating the complex resulting from step (b);
- d) determining the sequence of the mRNA in the isolated complex, thereby determining the identity of the gene.

14. The method of claim 13, wherein step (d) comprises steps of:

- i) synthesizing complementary DNA to the mRNA; and
- ii) performing sequence analysis of the complementary DNA to determine the sequence of the mRNA.

15. A gene identified by the method of claim 13.

16. The gene of claim 15 designated *MUM-1*.

17. The gene of claim 15 designated *MUM-2*.

18. The method of claim 13, wherein the gene identified comprises a nucleic acid encoding a MUM protein.

19. The method of claim 18, wherein the MUM protein is MUM-1.

20. The method of claim 18, wherein the MUM protein is MUM-2.

21. An isolated nucleic acid molecule encoding a MUM protein.

22. An isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule is a DNA molecule.

23. The isolated DNA molecule of claim 21, wherein the DNA molecule is a cDNA molecule.

24. The isolated DNA molecule of claim 21, wherein the DNA molecule is a cDNA molecule having the nucleotide sequence shown in Figure 5B (SEQ. ID NO ).

25. The isolated DNA molecule of claim 21, wherein the DNA molecule is genomic DNA molecule.

26. The isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule is an RNA molecule.
27. An isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule encodes a human MUM-1 protein.
28. An isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule encodes a human MUM-2 protein.
29. An isolated nucleic molecule of claim 31, wherein the human MUM-1 protein has substantially the same amino acid sequence as shown in Figure 5B (SEQ. ID NO ).
30. An isolated nucleic molecule of claim 31, wherein the human MUM-1 protein has the amino acid sequence as shown in Figure 5B (SEQ. ID NO ).
31. An isolated nucleic acid molecule of claim 21 operatively linked to a promoter of RNA transcription.
32. A vector comprising the nucleic acid molecule of claim 21.
33. A vector comprising the nucleic acid molecule of any claim 23.
34. A vector comprising the nucleic acid molecule of claim 25.
35. A vector of claim 36, wherein the vector is a plasmid.
36. The plasmid of claim 35, designated pcMUM1-1.6a (ATCC

Accession No.            ).

37. The plasmid of claim 35, designated pMUM1-2.4B/N (ATCC  
Accession No.            ).

5

38. The plasmid of claim 35, designated pMUM1-7.7B (ATCC  
Accession No.            ).

10

39. The plasmid of claim 35, designated pcMUM2-8 (ATCC  
Accession No.            ).

40. A host cell comprising the vector of claims 32.

15

41. The host cell of claim 40, wherein the cell is selected  
from a group consisting of a bacterial cell, a plant  
cell, and insect cell and a mammalian cell.

20

42. A nucleic acid probe comprising a nucleic acid molecule  
of at least 15 nucleotides capable of specifically  
hybridizing with a unique sequence included within the  
sequence of a nucleic acid molecule encoding a MUM  
protein.

25

43. A nucleic acid probe comprising a nucleic acid molecule  
of at least 15 nucleotides which is complementary to a  
sequence of the isolated nucleic acid molecule encoding  
a MUM protein.

30

44. The nucleic acid probe of either of claims 42 or 43,  
wherein the MUM protein is MUM-1.

45. The nucleic acid probe of either of claim 42 or 43,  
wherein the MUM protein is MUM-2.

46. A DNA probe of claim 44 or 45.

47. A RNA probe of claim 44 or 45.

5 48. A genomic DNA probe of claim 44 or 45.

49. A nucleic acid probe of claim 44 or 45 labeled with a detectable marker.

10 50. The nucleic acid probe of claim 49, wherein the detectable marker is selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

15 51. A nucleic acid probe of claim 44, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked to a nucleic acid sequence capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule  
20 of human chromosome 14.

52. A nucleic acid probe of claim 45, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked to a nucleic acid sequence capable of  
25 specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule of human chromosome 14.

53. A nucleic acid probe comprising a nucleic acid molecule  
30 of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule of claim 21 which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.



54. The nucleic acid probe of claim 53, wherein the isolated nucleic acid molecule encodes MUM-1.

5 55. The nucleic acid probe of claim 53, wherein the isolated nucleic acid molecule encodes MUM-2.

10 56. The nucleic acid probe of claim 54, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

15 57. The nucleic acid probe of claim 55, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

20 58. The nucleic acid probe of claim 56, wherein the specific break point comprises a portion of the t(6;14)(p25;q32) translocation.

25 59. The nucleic acid probe of claim 57, wherein the specific break point comprises a portion of a t(14;15) translocation.

30 60. The nucleic acid probe of either of claims 58 or 59 labeled with a detectable marker.

61. The nucleic acid probe of claim 60, wherein the detectable marker is selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

62. A method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1

protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein.

5      63. A method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein.

10

64. The method of claim 62, wherein the rearrangement of nucleic acid encoding MUM-1 protein is detected by contacting the nucleic acid from the sample with a MUM-1 probe under conditions permitting the MUM-1 probe to hybridize with the nucleic acid encoding MUM-1 protein from the sample, thereby detecting the rearrangement of nucleic acid encoding MUM-1 protein in the sample.

15

65. The method of claim 63, wherein the rearrangement of nucleic acid encoding MUM-2 protein is detected by contacting the nucleic acid from the sample with a MUM-2 probe under conditions permitting the MUM-2 probe to hybridize with the nucleic acid encoding MUM-2 protein from the sample, thereby detecting the rearrangement of nucleic acid encoding MUM-2 protein in the sample.

20

25

66. The method of claim 64, wherein the MUM-1 probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

30

35      67. The method of claim 65, wherein the MUM-2 probe

comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

68. The method of claim 66, wherein the nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

69. The method of claim 67, wherein the nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

70. The method of claim 68, wherein the specific break point comprises a portion of the t(6;14)(p25;q32) translocation.

71. The method of claim 69, wherein the specific break point comprises a portion of a t(14;15) translocation.

72. The method of claim 62, which comprises:

- a. obtaining DNA from the sample of the subject suffering from multiple myeloma;

- b. performing a restriction digest of the DNA with a panel of restriction enzymes;
- c. separating the resulting DNA fragments by size fractionation;
- d. contacting the resulting DNA fragments with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker;
- e. detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma;
- f. preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and
- g. comparing the detected band pattern specific to the DNA obtained from a sample of

subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

73. The method of claim 63, which comprises:

- a. obtaining DNA from the sample of the subject suffering from multiple myeloma;
- b. performing a restriction digest of the DNA with a panel of restriction enzymes;
- c. separating the resulting DNA fragments by size fractionation;
- d. contacting the resulting DNA fragments with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker;
- e. detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2

protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma;

f. preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and

g. comparing the detected band pattern specific to the DNA obtained from a sample of subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

74. The method of claim 72 or 73, wherein the size fractionation in step (c) is effected by a polyacrylamide or agarose gel.

75. The method of claim 72 or 73, wherein the detectable marker is radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

76. A method of claim 62 which comprises:

a. obtaining RNA from the sample of the subject suffering from multiple myeloma;

- b. separating the RNA sample by size fractionation;
- c. contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker;
- d. detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma;
- f. preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and
- g. comparing the detected band pattern specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

77. A method of claim 63 which comprises:

- a. obtaining RNA from the sample of the subject

suffering from multiple myeloma;

b. separating the RNA sample by size fractionation;

5

c. contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 15 and labeled with a detectable marker;

10

15

d. detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma;

20

f. preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and

25

g. comparing the detected band pattern specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

30

35 78. The method of claim 76 or 77, wherein the size



fractionation in step (c) is effected by a polyacrylamide or agarose gel.

- 5 79. The method of claim 76 or 77, wherein the detectable marker is radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.
- 10 80. The method of either of claim 72 or 76, wherein multiple myeloma associated with the expression of a specific human MUM-1 is diagnosed.
- 15 81. The method of either of claim 73 or 77, wherein multiple myeloma associated with the expression of a specific human MUM-2 is diagnosed.
- 20 82. An antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule.
- 25 83. An antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule.
- 30 84. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the cDNA molecule of claim 23.
85. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the genomic DNA molecule of claim 29.
86. An antisense oligonucleotide having a sequence capable

of specifically hybridizing to the RNA molecule of claim 30.

87. An purified MUM protein.

5

88. A purified MUM protein, wherein the MUM protein is MUM-1 protein.

89. A purified human MUM-1 protein of claim 88.

10

90. An antibody directed to a purified MUM-1 protein.

91. An antibody capable of specifically recognizing MUM-1 protein.

15

92. An antibody of claim 91, wherein the MUM-1 protein is a human MUM-1 protein.

93. A purified MUM protein, wherein the MUM protein is MUM-2 protein.

20

94. A purified human MUM-2 protein of claim 93.

95. An antibody directed to a purified MUM-2 protein.

25

96. An antibody capable of specifically recognizing a MUM-2 protein.

97. An antibody of claim 96, wherein the MUM-2 protein is a human MUM-2 protein.

30

98. An monoclonal antibody of any one of claims 90, 91 and 92.

99. An monoclonal antibody of any one of claims 95, 96, and 97.

5 100. A pharmaceutical composition comprising an amount of the oligonucleotide of any one of claims 82, 84, 85 and 81 effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

10 101. A pharmaceutical composition comprising an amount of the oligonucleotide of any one of claims 83, 84, 85 and 81 effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

15

IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

Abstract of the Invention

5

10

15


20

25

30

This invention provides a method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of: (a) obtaining a DNA sample from the subject suffering from multiple myeloma; (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample; (c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction; (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene; (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones from step (d); (f) hybridizing said fluorescent probes with metaphase chromosomes; and (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma. This invention also provides the identified gene altered by a chromosomal breakpoint and various uses thereof.

FIG. 1




14q32 (IgH) germline

FIG. 2A


5kb



der.6  
(λSKB-4a)



der.14  
(λSKS-3)



6p25 germline




FIG. 2B

↓

.6 TTTTCTCTACAGTCACCTCCCTGTTACCAAGATAATCACATAAGTCCAGTTTACTTACAAACAAGTTAGT  
 |||||  
 .6 TGGGCTCGGCC-TGGTGGGGCAGCCACAGCGGGACGC-AGTAGTGAAAGTCCAGTTTACTTACAAACAAGTTAGT  
 |||||  
 14 TGGGCTCGGCCTTGGTGGGCAGCCACAGCGGGACGCAAGTAGTGAGGGCACTCAGAACGCCACTCAGCCCCGACAG  
 4400 4410 4420 4430 4440 4450 4460 4470

↓

.6 TATTAGAGGAAACTAAAACTTCAGGATTCAGTCCAGATAATTTTAAAACTCTAAACAATGGACAGGGCTAGAAT  
 |||||  
 .14 TATTAGAGGAAACTAAAACTTCAGGATTCAGCAGGGCATGAGGAGGCAGCTCCTCACCCCTTCTCTTTGTAC  
 |||||  
 .14 GGCACTCAGAACGCCACTCAGCCCCGACAGGCAGGGCACGAGGAGGCAGCTCCTCACCCCTTCTCTTTGT--  
 4450 4460 4470 4480 4490 4500 4510

FIG. 3A



FIG. 3B




$\lambda$ MUM-3




5/21

FIG. 4A



6/21

FIG. 4B



7/21

FIG. 4C





FIG. 5A





1151	AGAGGGGCGT	GGTCTCTG	ATGGCCCCCG	ACGGGCTCTA	TGCGAAAAAC
	R G V	V L W	M A P D	G L Y	A K R
1201	CTGTGCCAGA	GCACGATCTA	CTGGGACGGG	CCCCTGGCGC	TGTGCAAC
	L C Q S	T I Y	W D G	P L A L	C N
1251	CCGGCCCAAC	AAACTGGAGA	GAGACCAGAC	CTGCAAGCTC	TTTGACAC
	R P N	K L E R	D Q T	C K L	F D T
1301	AGCAGTTCTT	GTCAGAGCTG	CAAGCGTTTG	CTCACCACGG	CCGCTCCC
	Q F L	S E L	Q A F A	H H G	R S L
1351	CCAAGATTCC	AGGTGACTCT	ATGCTTTGGA	GAGGAGTTTC	CAGACCCT
	P R F Q	V T L	C F G	E E F P	D P
1401	GAGGCAAAGA	AAGCTCATCA	CAGCTCACGT	AGAACCTCTG	CTAGCCAG
	R Q R	K L I T	A H V	E P L	L A R
1451	AACTATATTA	TTTTGCTCAA	CAAAACAGTG	GACATTTCTT	GAGGGGCT
	L Y Y	F A Q	Q N S G	H F L	R G Y
1501	GATTTACCAG	AACACATCAG	CAATCCAGAA	GATTACCACA	GATCTATC
	D L P E	H I S	N P E	D Y H R	S I
1551	CCATTCCCTCT	ATTCAAGAAT	GAAAAATGTC	AAGATGAGTG	GTTTTCTT
	H S S	I Q E *			
1601	TCCTTTTTTTT	TTTTTTTTTTT	TTTGATACGG	AGATACGGGG	TCTTGCTC
1651	TCTCCCAGGC	TGGAGTGCAG	TGACACAATC	TCAGCTCACT	GTGACCTC
1701	CCTCCTGGGT	TCAAGAGACT	CTCCTGCCTC	AGCCTCCCTG	GTAGCTGG
1751	TTACAGGTGT	GAGCCACTGC	ACCCACCCAA	GACAAGTGAT	TTTCATTG
1801	AATATTTGAC	TTTAGTGAAA	GCGTCCAATT	GACTGCCCTC	TTACTGTT
1851	GAGGAACTCA	GAAGTGGAGA	TTTCAGTTCA	GCGGTTGAGG	AGAATTGC
1901	CGAGACAAGC	ATGGAAAATC	AGTGACATCT	GATTGGCAGA	TGAGCTTA
1951	TCAAAAGGAA	GGGTGGCTTT	GCATTTTCTT	GTGTTCTGTA	GACTGCCA
2001	ATTGATGATC	ACTGTGAAAA	TTGACCAAGT	GATGTGTTTA	CATTTACT
2051	AATGCGCTCT	TTAATTTGTT	GTAGATTAGG	TCTTGCTGGA	AGACAGAG
2101	AACTTGCCTT	TCAGTATTGA	CACTGACTAG	AGTGATGACT	GCTTGTAG
2151	ATGTCTGTGC	CATTTCTCAG	GGAAGTAAGA	TGTAAATTGA	AGAAGCCT
2201	CACGTAAAAG	AAATGTATTA	ATGTATGTAG	GAGCTGCAGT	TCTTGTGG
2251	GACACTTGCT	GAGTGAAGGA	AATGAATCTT	TGACTGAAGC	CGTGCCCTG
2301	GCCTTGGGGA	GGCCCATCCC	CCACCTGCCA	GCGGTTTCTT	GGTGTGGG
2351	CCTCTGCCCC	ACCCTCCTTC	CCATTGGCTT	TCTCTCCTTG	GCCTTTCC
2401	GAAGCCAGTT	AGTAAACTTC	CTATTTTCTT	GAGTCAAAAA	ACATGAGC
2451	TACTCTTGGA	TGGGACATTT	TTGTCTGTCC	TACAATCTAG	TAATGTCT
2501	GTAATGGTTA	AGTTTTCTTG	TTTCTGCATC	TTTTTGACCC	TCATTCTT
2551	GAGATGCTAA	AATTCTTCGC	ATAAAGAAGA	AGAAATTAAG	GAACATAA
2601	CTTAATACTT	GAAGTGTGTC	CCTTCTGTCC	AAGTACTTAA	CTATCTGT
2651	CCTTCCTCTG	TGCCACGCTC	CTCTGTTTGT	TTGGCTGTCC	AGCGATCA
2701	CATGGCGACA	CTAAAGGAGG	AGGAGCCGGG	GAATCCCAGG	CTGGAGAG
2751	CTGCCAGGAC	CCACCACTGG	AAGCAGGATG	GAGCTGACTA	CGGAACCTG
2801	CACTCAGTGG	GCTGTTTCTG	CTTATTTTCAT	CTGTTCTATG	CTTCCTCG
2851	CCAATTATAG	TTTGACAGGG	CCTTAAAATT	ACTTGCTTTT	TTCCAAAT
2901	TTCTATTTAT	AGAAATCCCA	AAGACCTCCA	CTTGCTTAAG	TATACCTA
2951	ACTTACATTT	TTGTGGTTTT	GAGAAAGTAC	AGCAGTAGAC	TGGGGCGT

FIG. 5B-3

3051	TCAGCAGAAG	ATTGCGTTAG	CTCTCAAATG	TGTGTTCCCTG	CTTTTCTA
3101	GGATATTTTA	AATTCATTCA	ACAAGCACCT	AGTAAGTGCC	TGCTGTAT
3151	CTACATTACA	CAGTTCAGCC	TTTATCAAGC	TTAGTGAGCA	GTGAGCAC
3201	AAACATTATT	TTTTAATGTT	TAAAAAGTTT	CTAATATTAA	AGTCAGAA
3251	TTAATACAAT	TAATATTAAAT	ATTAACTACA	GAAAAGACAA	ACAGTAGA
3301	ACAGCAAAAA	AATAAAAAGG	ATCTCCTTTT	TTCCCAGCCC	AAATTCTC
3351	CTCTAAAAGT	GTCCACAAGA	AGGGGTGTTT	ATTCTTCCAA	CACATTTT
3401	TTTTCTGTAA	ATATACATAA	ACTTAAAAAG	AAAACCTCAT	GGAGTCAT
3451	TGCACACACT	TTTCATGCAG	TGCTCTTTGT	AGCTAAACAG	TGAAGATT
3501	CCTCGTTCTG	CTCAGAGGCC	TTGCTGTGGA	GCTCCACTGC	CATGTACC
3551	GTAGGGTTTG	ACATTTTCAT	AGCCATGCAA	CATGGATATG	TATTGGGC
3601	CAGACTGTGT	TTCGTGAAC	GCAGTGATGT	ATACATCTTA	TAGATGCA
3651	GTATTTTGGG	GTATATTATC	CTAAGGGAAG	ATAAAGATGA	TATTAAGA
3701	TGCTGTTTCA	CGGGGCCCTT	ACCTGTGACC	CTCTTTGCTG	AAGAATAT
3751	AACCCACAC	AGCACTTCAA	AGAAGCTGTC	TTGGAAGTCT	GTCTCAGG
3801	CACCCTGTCT	TCTTAATTCT	CCAAGCGGAT	GCTCCATTTC	AATTGCTT
3851	TGACTTCTTC	TTCTTTGTTT	TTTTAAATAT	TATGCTGCTT	TAACAGTG
3901	GCTGAATTTT	CTGGAAAATG	CTTCTTGCTT	GGGGCCACTA	CCTCCTTT
3951	TATCTTTACA	TCTATGTGTA	TGTTGACTTT	TTAAAATTCT	GAGTGATC
4001	GGGTATGACC	TAGGGAATGA	ACTAGCTATG	GAAATAACTC	AGGGTTAG
4051	ATCCTAGCAC	TTGTCTCAGG	ACTCTGAAAA	GGAACGGCTT	CCTCATTC
4101	TGTCTTGATA	AAGTGGAATT	GGCAAAC TAG	AATTTAGTTT	GTACTCAG
4151	GACAGTGCTG	TTGAAGATTT	GAGGACTTGT	TAAAGAGCAC	TGGGTCAT
4201	GGAAAAAATG	TATGTGTCTC	CCCAGGTGCA	TTTTCTTGGT	TTATGTCT
4251	TTCTTGAGAT	TTTGTATATT	TAGGAAAACC	TCAAGCAGTA	ATTAATAT
4301	CCTGGAACAC	TATAGAGAAC	CAAGTGACCG	ACTCATTTAC	AACTGAAAC
4351	TAGGAAGCCC	CTGAGTCCTG	AGCGAAAACA	GGAGAGTTAG	TGCCCCTAC
4401	GAAAACCCAG	CTAGACTATT	GGGTATGAAC	TAAAAAGAGA	CTGTGCCAT
4451	GTGAGAAAAA	TGTAAAATCC	TACAGTGGA	TGAGCAGCCC	TTACAGTG
4501	GTTACCACCA	AGGGCAGGTA	GGTATTAGTG	TTTGAAAAAG	CTGGTCTTT
4551	AGCGAGGGCA	TAAATACAGC	TAGCCCCAGG	GGTGGAACAA	CTGTGGGAC
4601	CTTGGGTACT	CGCACCTCTT	GGCTTTGTTG	ATGCTCCGCC	AGGAAGGCC
4651	CTTGTGTGTG	CGTGT CAGTT	ACTTTTTTTAG	TAACAATTCA	GATCCAGTC
4701	AAACTTCCGT	TCATTGCTCT	CCAGTCACAT	GCCCCCACTT	CCCCACAGG
4751	GAAAGTTTTT	CTGAAGTGTT	GGGATTGGTT	AAGGTCTTTA	TTTGTATTA
4801	GTATCTCCCC	AAGTCCTCTG	TGGCCAGCTG	CATCTGTCTG	AATGGTGCG
4851	GAAGGCTCTC	AGACCTTACA	CACCATTTTG	TAAGTTATGT	TTTACATGC
4901	CCGTTTTTTGA	GACTGATCTC	GATGCAGGTG	GATCTCCTTG	AGATCCTGA
4951	AGCCTGTTAC	AGGAATGAAG	TAAAGGTCAG	TTTTTTTTTG	ATTGATTTT
5001	ACAGCTTTGA	GGAACATGCA	TAAGAAATGT	AGCTGAAGTA	GAGGGGACG
5051	GAGAGAAGGG	CCAGGCCGGC	AGGCCAACCC	TCCTCCAATG	GAAATTCCC
5101	TGTTGCTTCA	AACTGAGACA	GATGGGACTT	AACAGGCAAT	GGGGTCCAC
5151	TCCCCCTCTT	CAGCATCCCC	CGTACC		

A

FIG. 6A

MUM-1 (23-72)	KLRQWLIDQI	DSGKYPGLMW	ENKESIFRI	PWKHAGKQDY	NREDDAALEK
LSIRF (23-72)	KLRQWLIDQI	DSGKYPGLMW	ENKESVFR	PWKHAGKQDY	NREDDAALEK
IRF-1 (7-56)	RMRPWLEQI	NSNQIPGLIW	INKEEMIFQI	PWKHAKHGW	DINKDACLER
IRF-2 (7-56)	RMRPWLEQI	NSNTIPGLKW	INKEKKIFQI	PWKHAKHGW	DVEKDAPLER
ICSBP (9-60)	RLRQWLIEQI	DSSMYPGLIW	ENKESMFRI	PWKHAGKQDY	NQLEVDASIEK
ISGF3Y (11-60)	KLRNWWVEQV	ESGQFPGVCW	DDTAKTMR	PWKHAGKQDF	REDQDAAFK
IRF-3 (7-55)	RILPWLVSQL	DLGQLEGVAM	VNKSRTFR	PWKHGTROD.	AQQEDFGIEQ
MUM-1 (73-122)	AWALEKKGKFR	EGIDKPDPPPT	MKTRLRCAIN	KSNDDEELVE	RSQLDISDPY
LSIRF (73-122)	AWALEKKGKFR	EGIDKPDPPPT	MKTRLRCAIN	KSNDDEELVE	RSQLDISDPY
IRF-1 (57-106)	SWAIHTGRYK	AGEKEPDPKTI	MKANFRCAIN	SLPDIIEEVKQ	QGRNKGSAAV
IRF-2 (57-106)	NWAIHTGKHQ	PGMDKPDPPKT	MKANFRCAIN	SLPDIIEEVKQ	KSIIKKGNAF
ICSBP (59-107)	AWALEKKGKFR	EGDKAEPAT	MKTRLRCAIN	KSPDEEEVTD	RSQLDISEPY
ISGF3Y (61-109)	AWALEKKGKYK	EGDTGCPAV	MKTRLRCAIN	KSSEKEVPE	RGRMDVAEPY
IRF-3 (56-104)	AWAEATGAYV	PGRDKPDLPPT	MKNFRSALN	RKEGLRLAED	RSKDPHDPH
MUM-1 (123-130)	KVYRIVPE				
LSIRF (123-130)	KVYRIVPE				
IRF-1 (107-114)	KVYRMIPP				
IRF-2 (107-114)	KVYRMIPPL				
ICSBP (108-115)	KVYRIVPE				
ISGF3Y (110-117)	KVYQLIPP				
IRF-3 (105-112)	KIYEFVNS				



FIG. 6B

MUM-1 (327-372)	KRLCOSTIYV	DGPLAL....	CNDRPNKIER	DOTCKLFDTO	QFLSEDA
LSIRF (327-372)	KRLCOSRIYV	DGPLAL....	CSDRPNKIER	DOTCKLFDTO	QFLSEDA
ICSBP (289-334)	KRLCOGRVFC	SGNAV....	CKGRPNKIER	DEVVQVETS	QFREFQY
ISGF3Y (290-335)	QRLCPIPIST	NAPQAP....	PGCPHLIPS	NECVELRPA	YACRDVRYF
IRF-3 (284-333)	QRLGHCHTYV	AVSEELLPNS	GHGPDGEVPK	KEGGVBLG	PLIVDITIT
MUM-1 (373-421)	HHGRSLPRFQ	VTLQFGELFP	DPQRQR.KH	TAHVPLLAN	QIYYFAQONS
LSIRF (373-421)	HHGRPAPRFQ	VTLQFGELFP	DPQRQR.KH	TAHVPLLAN	QIYYFAQONT
ICSBP (335-384)	NSQGRLPDGR	VVICFGELFP	DMAPLRSKH	LVQIIPOLYR	QIAEEDGKSC
ISGF3Y (336-385)	QGLGPPPKFQ	VVINMESH	GSSHTPONAI	TVKMDOAFK	YILEQTPEQQ
IRF-3 (334-383)	EGSGRSRYA	LWFVGSMP	QDQPWTKRIV	MVKVPTCLR	AVEMRVGG



FIG. 8




FIG. 9A



FIG. 9B




FIG. 10

*MUM2* Transcripts detected in MM/PCL Cell Lines




FIG. 11A

## Physiological IgH gene rearrangement






FIG. 11B

## Chromosomal translocation occurring in switch region



## MUM1 cDNA



Sc; SacII, P; *PstI*, H; *HindIII*, S; *SacI*, A; *ApaI*

cDNA inserts is cloned into EcoRI / BamHI site of the pBluescript KS+  
Bacteria strain used is DH5 $\alpha$  cells. pcMUM1-1.6a contains full length open  
reading frame of nt.217-1572.





FIG. 12 A-B



# Breakpoint Cloning of the U-266 Cell Line



pMUM2-8 has a 22.0kb insert in BamHI site of pBluescript KS+.

FIG. 13

[illegible]

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

<u>Provisional Application No.</u>	<u>Filing Date</u>	<u>Status</u>
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States Application(s), or Section 365(c) of any PCT International Application(s) designating the United States listed below. Insofar as this application discloses and claims subject matter in addition to that disclosed in any such prior Application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56, which became available between the filing date(s) of such prior Application(s) and the national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

And I hereby appoint

John P. White (Reg. No. 28,678); Thomas F. Moran (Reg. No. 16,579); Norman H. Zivin (Reg. No. 25,385); Ivan S. Kavrukov (Reg. No. 25,161); Christopher C. Dunham (Reg. No. 22,031); Thomas G. Carulli (Reg. No. 30,616); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Richard S. Milner (Reg. No. 33,970); Albert Wai-Kit Chan (Reg. No. 36,479); Kristina L. Konstas (Reg. No. 37,864); Mary Anne P. Tanner (Reg. No. 40,197); Timothy X. Witkowski (Reg. No. 40,232); and Mary Catherine DiNunzio (Reg. No. 37,306)

and each of them, all c/o Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, New York 10036, my attorneys, each with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent, to transact all business in the Patent and Trademark Office connected therewith and to file any International Applications which are based thereon under the provisions of the Patent Cooperation Treaty.

Please address all communications, and direct all telephone calls, regarding this application to:

John P. White \_\_\_\_\_ Reg. No. 28,678  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10036  
Tel. (212) 278-0400

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or  
first joint inventor \_\_\_\_\_ Riccardo Dalla-Favera

Inventor's signature \_\_\_\_\_ *R. Dalla-Favera*

Citizenship \_\_\_\_\_ Italy \_\_\_\_\_ Date of signature \_\_\_\_\_ 6/6/96

Residence \_\_\_\_\_ 445 Riverside Drive, #122, New York, New York 10025

Post Office Address \_\_\_\_\_ same as residence

Full name of joint  
inventor (if any) \_\_\_\_\_

Inventor's signature \_\_\_\_\_

Citizenship \_\_\_\_\_ Date of signature \_\_\_\_\_

Residence \_\_\_\_\_

Post Office Address \_\_\_\_\_

Full name of joint  
inventor (if any) \_\_\_\_\_

Inventor's signature \_\_\_\_\_

Citizenship \_\_\_\_\_ Date of signature \_\_\_\_\_

Residence \_\_\_\_\_

Post Office Address \_\_\_\_\_

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Riccardo Dalla-Favera  
U.S. Serial No.: Not Yet Known (Continuation of U.S. Serial  
No. 08/654,482, filed May 28, 1996)  
Filed : Herewith  
For : IDENTIFICATION OF GENES ALTERED IN MULTIPLE  
MYELOMA

1185 Avenue of the Americas  
New York, New York 10036  
June 1, 2000

Assistant Commissioner for Patents  
Washington, D.C. 20231

Box: Patent Application

SIR:

ASSOCIATE POWER OF ATTORNEY


Please recognize Jay H. Maioli (Reg. No. 27,213); William E. Pelton  
(Reg. No. 25,702); Wendy E. Miller (Reg. No. 35,615); Robert T.  
Maldonado (Reg. 38,232); Paul Teng (40,837); Richard F. Jaworski  
(Reg. No. 33,515); Elizabeth M. Wieckowski (Reg. No. 42,226);  
Pedro C. Fernandez (Reg. No. 41,741); Gary J. Gershik (Reg. No.  
39,992); Jane M. Love (Reg. No. 42,812); Spencer H. Schneider (Reg.  
No. 45,923); and Raymond A. Diperna (Reg. No. 44,063) as associate  
attorney in connection with the above-identified application.

Applicants: Riccardo Dalla-Favera  
U.S. Serial No.: Not Yet Known  
Filed: Herewith  
Page 2

Plase direct correspondence to the attorney of record at the following address:

John P. White  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10016  
(212) 278-0400

Respectfully submitted,

A handwritten signature in dark ink, appearing to read "John P. White", is written over a horizontal line.

John P. White  
Registration No. 28,678  
Attorney for Applicant  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10016  
(212) 278-0400